

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
22 February 2001 (22.02.2001)

PCT

(10) International Publication Number
WO 01/12186 A1

(51) International Patent Classification⁷: A61K 31/40,
31/4025, 31/445, C07D 401/06, 401/08, 401/10, 401/12

02493 (US). CORNEBISE, Mark [US/US]; 22 Gilbert
Street, Watertown, MA 02472 (US). PETTER, Russell
[US/US]; 343 Hudson Road, Stow, MA 01775 (US).

(21) International Application Number: PCT/US00/22285

(74) Agent: MYERS, Louis; Fish & Richardson P.C., 225
Franklin Street, Boston, MA 02110-2804 (US).

(22) International Filing Date: 14 August 2000 (14.08.2000)

(25) Filing Language: English

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(26) Publication Language: English

(30) Priority Data:
60/148,845 13 August 1999 (13.08.1999) US

(63) Related by continuation (CON) or continuation-in-part
(CIP) to earlier application:
US 60/148,845 (CIP)
Filed on 13 August 1999 (13.08.1999)

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): BIO-
GEN, INC. [US/US]; 14 Cambridge Center, Cambridge,
MA 02142 (US).

Published:

— With international search report.

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): LEE, Wen-Cherng
[—/US]; 192 Spring Street, Lexington, MA 02421 (US).
SCOTT, Daniel [US/US]; 42 Bradyll Road, Weston, MA

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*



WO 01/12186 A1

(54) Title: CELL ADHESION INHIBITORS

(57) Abstract: A cell adhesion inhibitor of the general formula: R³-L'-R¹ is disclosed. An inhibitor of the present invention interacts with VLA-4 molecules and inhibits VLA-4 dependent cell adhesion. Also disclosed are methods for preparing and using such a cell adhesion inhibitor, as well as pharmaceutical compositions containing the same.

CELL ADHESION INHIBITORS

5

BACKGROUND

Cell adhesion is a process by which cells associate with each other, migrate towards a specific target or localize within the extra-cellular matrix. As such, cell adhesion constitutes one of the fundamental mechanisms underlying numerous biological phenomena. For example, cell adhesion is responsible for the adhesion of hematopoietic cells to endothelial cells and the subsequent migration of those hemopoietic cells out of blood vessels and to the site of injury. As such, cell adhesion plays a role in pathologies such as inflammation and immune reactions in mammals.

Investigations into the molecular basis for cell adhesion have revealed that various cell-surface macromolecules -- collectively known as cell adhesion molecules or receptors -- mediate cell-cell and cell-matrix interactions. For example, proteins of the superfamily called "integrins" are key mediators in adhesive interactions between hematopoietic cells and their microenvironment (M.E. Hemler, "VLA Proteins in the Integrin Family: Structures, Functions, and Their Role on Leukocytes.", Ann. Rev. Immunol., 8, p. 365 (1990)). Integrins are non-covalent heterodimeric complexes consisting of two subunits called α and β . There are at least 12 different α subunits ($\alpha 1$ - $\alpha 6$, α -L, α -M, α -X, α -IIB, α -V and α -E) and at least 9 different β ($\beta 1$ - $\beta 9$) subunits. Based on the type of its α and β subunit components, each integrin molecule is categorized into a subfamily.

$\alpha 4\beta 1$ integrin, also known as very late antigen-4 ("VLA-4"), CD49d/CD29, is a leukocyte cell surface receptor that participates in a wide variety of both cell-cell and cell-matrix adhesive interactions (M.E. Hemler, Ann. Rev. Immunol., 8, p. 365 (1990)). It serves as a receptor for the cytokine-inducible endothelial cell surface protein, vascular cell adhesion molecule-1 ("VCAM-1"), as well as to the extracellular matrix protein fibronectin ("FN") (Ruegg et al., J. Cell Biol., 177, p. 179 (1991); Wayner et al., J. Cell Biol., 105, p. 1873 (1987); Kramer et al., J. Biol. Chem., 264, p. 4684 (1989); Gehlsen et al. Science, 24, p.

1228 (1988)). Anti-VLA4 monoclonal antibodies ("mAb's") have been shown to inhibit VLA4-dependent adhesive interactions both *in vitro* and *in vivo* (Ferguson et al. *Proc. Natl. Acad. Sci.*, 88, p. 8072 (1991); Ferguson et al., *J. Immunol.*, 150, p. 1172 (1993)). Results of *in vivo* experiments suggest that this inhibition of VLA-4-dependent cell adhesion may prevent or inhibit several inflammatory and autoimmune pathologies (R. L. Lobb et al., "The Pathophysiologic Role of $\alpha 4$ Integrins In Vivo", *J. Clin. Invest.*, 94, pp. 1722-28 (1994)).

Despite these advances, there remains a need for small, specific inhibitors of VLA-4-dependent cell adhesion. Ideally, such inhibitors may be orally administered. Such compounds would provide useful agents for treatment, prevention or suppression of various pathologies mediated by cell adhesion and VLA-4 binding.

SUMMARY

The present invention relates to novel non-peptidic compounds that specifically inhibit the binding of ligands to VLA-4. These compounds are useful for inhibition, prevention and suppression of VLA-4-mediated cell adhesion and pathologies associated with that adhesion, such as inflammation and immune reactions. The compounds of this invention may be used alone or in combination with other therapeutic or prophylactic agents to inhibit, prevent or suppress cell adhesion. This invention also provides pharmaceutical compositions containing the compounds of this invention and methods of using the compounds and compositions of the invention for inhibition of cell adhesion.

According to one embodiment of this invention, these novel compounds, compositions and methods are advantageously used to treat inflammatory and immune diseases. The present invention also provides methods for preparing the compounds of this invention and intermediates therefor.

An aspect of this invention relates to cell adhesion inhibitors of formula (I):




R^1 is H, C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, Cy, Cy- C_{1-10} alkyl, Cy- C_{1-10} alkenyl, or Cy- C_{1-10} alkynyl.

L' is a hydrocarbon linker moiety having 1-5 carbon chain atoms and is (i) optionally interrupted by, or terminally attached to, one or more (e.g., 1, 2, or 3) of the following groups: -C(O)-, -O-C(O)-, -C(O)-O-, -C(O)-NR^c-, -NR^c-C(O)-, -NR^c-C(O)-NR^d-, -NR^c-

C(O)-O-, -O-C(O)-NR^c-, -S(O)_m-, -SO₂-NR^c-, -NR^c-SO₂-, -NR^c-C(NR^m)-, -O-, -NR^c-, and -Cy; or (ii) optionally substituted with one or more substituents independently selected from R^b.

L is a hydrocarbon linker moiety having 1-14 carbon chain atoms and is (i) optionally interrupted by, or terminally attached to, one or more (e.g., 1-5, 1-4, or 1-3) of the following groups: -C(O)-, -O-C(O)-, -C(O)-O-, -C(O)-NR^c-, -NR^c-C(O)-, -NR^c-C(O)-NR^d-, -NR^c-C(O)-O-, -O-C(O)-NR^c-, -S(O)_m-, -SO₂-NR^c-, -NR^c-SO₂-, -O-, -NR^c-, and Cy; or (ii) optionally substituted with one or more substituents independently selected from R^b.

R³ is alkyl, alkenyl, alkynyl, cycloalkyl, aryl-fused cycloalkyl, cycloalkenyl, aryl, aralkyl, aryl-substituted alkenyl or alkynyl, cycloalkyl-substituted alkyl, cycloalkenyl-substituted cycloalkyl, biaryl, alkenoxy, alkynoxy, aralkoxy, aryl-substituted alkenoxy, aryl-substituted alkynoxy, alkylamino, alkenylamino, alkynylamino, aryl-substituted alkylamino, aryl-substituted alkenylamino, aryl-substituted alkynylamino, aryloxy, arylamino, heterocyclyl, heterocyclyl-substituted alkyl, heterocyclyl-substituted amino, carboxyalkyl substituted aralkyl, or oxocarbocyclyl-fused aryl; or R³ is a moiety of formula (i):



Y⁵ is -CO-, -O-CO-, -SO₂- or -PO₂-.

Each of R⁴ and R⁶, independently, is alkyl, alkenyl, alkynyl, cycloalkyl, aryl-fused cycloalkyl, cycloalkenyl, aryl, aralkyl, aryl-substituted alkenyl or alkynyl, cycloalkyl-substituted alkyl, cycloalkenyl-substituted cycloalkyl, biaryl, alkenoxy, alkynoxy, aralkoxy, aryl-substituted alkenoxy, aryl-substituted alkynoxy, alkylamino, alkenylamino, alkynylamino, aryl-substituted alkylamino, aryl-substituted alkenylamino, aryl-substituted alkynylamino, aryloxy, arylamino, heterocyclyl, heterocyclyl-substituted alkyl, heterocyclyl-substituted amino, carboxyalkyl substituted aralkyl, oxocarbocyclyl-fused aryl, or an amino acid side chain selected from the group consisting of arginine, asparagine, glutamine, S-methyl cysteine, methionine and corresponding sulfoxide and sulfone derivatives thereof, cyclohexylalanine, leucine, isoleucine, allo-isoleucine, tert-leucine, norleucine, phenylalanine, phenylglycine, tyrosine, tryptophan, proline, alanine, ornithine, histidine,

glutamine, norvaline, valine, threonine, serine, beta-cyanoalanine, 2-aminobutyric acid and allothreonine.

R^5 is hydrogen, aryl, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, or aryl-substituted alkyl. Note that R^5 and R^6 may be taken together with the atoms to which they
5 are attached to form a heterocycle of 5 to 7 members.

Each of the above-stated Cy represents cycloalkyl, cycloalkenyl, heterocyclyl, aryl, or heteroaryl. Each of the above-stated alkyl, alkenyl and alkynyl is optionally substituted with one to four substituents independently selected from R^a . Further, each of the above-stated cycloalkyl, cycloalkenyl, heterocyclyl, aryl, and heteroaryl is optionally substituted with one
10 to four substituents independently selected from R^b .

R^a is selected from the group consisting of: Cy (which is optionally substituted with one to four substituents independently selected from R^b), $-OR^c$, $-NO_2$, -halogen, $-S(O)_mR^c$, $-SR^c$, $-S(O)_2OR^c$, $-S(O)_2NR^cR^d$, $-NR^cR^d$, $-O(CR^cR^f)_nNR^cR^d$, $-C(O)R^d$, $-CO_2R^c$, $-P(O)(OR^c)(OR^d)$, $-P(O)(R^c)(OR^d)$, $-S(O)_mOR^c$, $-C(O)NR^cR^j$, $-CO_2(CR^cR^f)_nCONR^cR^d$,
15 $OC(O)R^c$, $-CN$, $-NR^cC(O)R^d$, $-OC(O)NR^cR^d$, $-NR^cC(O)OR^d$, $-NR^cC(O)NR^dR^e$, $-CR^c(NOR^d)$, $-CF_3$, $-OCF_3$, and oxo.

R^b is a group selected from R^a , C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, aryl- C_{1-10} alkyl, and heteroaryl- C_{1-10} alkyl; wherein each of alkyl, alkenyl, alkynyl, aryl, and heteroaryl is optionally substituted with a group independently selected from R^e .

20 Each of R^c , R^d , R^e , and R^f , independently, is selected from H, C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, Cy, and Cy- C_{1-10} alkyl; wherein each of alkyl, alkenyl, alkynyl and Cy is optionally substituted with one to four substituents independently selected from R^e .

R^e is halogen, amino (including $-NH_2$, (mono- or di-)alkylamino, (mono- or di-) alkenylamino, (mono- or di-)alkynylamino, (mono- or di-)cycloalkylamino, (mono- or di-) cycloalkenylamino, (mono- or di-)heterocyclylamino, (mono- or di-)arylamino, and (mono- or di-)heteroarylamino), carboxy, $-COO-C_{1-4}$ alkyl, $-P(O)(OH)_2$, $-P(O)(OH)(O-C_{1-4}$ alkyl), $-P(O)(C_{1-4}$ alkyl) $_2$, $-P(O)(OH)(C_{1-4}$ alkyl), $-P(O)(O-C_{1-4}$ alkyl)(C_{1-4} alkyl), $-SO_2-C_{1-4}$ alkyl, $-CO-NH_2$, $-CO-NH(C_{1-4}$ alkyl), $-CO-N(C_{1-4}$ alkyl) $_2$, $-C_{1-4}$ alkyl, $-C_{1-4}$ alkoxy, aryl, aryl- C_{1-4} alkoxy, hydroxy, CF_3 , and aryloxy.

30 R^m is H, C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, Cy, Cy- C_{1-10} alkyl, C_{1-10} acyl, C_{1-10} alkyl-sulfonyl, or C_{1-10} alkoxy.

R^j is H, C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, cyano, aryl, aryl- C_{1-10} alkyl, heteroaryl, heteroaryl- C_{1-10} alkyl, or $-SO_2R^k$ (with R^k being C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, or aryl).


R^c and R^d can be taken together with the atoms to which they are attached and optionally form a heterocyclic ring of 5 to 7 members that contains 0-2 additional heteroatoms independently selected from O, N and S. Similarly, R^e and R^f can be taken together with the atoms to which they are attached optionally form a ring of 5 to 7 members that contains 0-2 additional heteroatoms independently selected from O, S and N.

m is 0, 1, or 2; and n is an integer from 1 to 10.

Note that when L is saturated (e.g., a C_{1-4} alkylene chain) and has 1-4 carbon chain atoms, L must contain a heteroatom selected from O, S, and N; or R^3 must contain the moiety o-methylphenyl-ureido-phenyl- CH_2 -; or R^1 must contain only one cyclic group (e.g., cycloalkyl, cycloalkenyl, heterocyclyl, aryl, or heteroaryl).

In one embodiment, the compounds of this invention contain R^1 with the formula: $Z^1-L^a-Z^2$, wherein Z^1 is cycloalkyl, cycloalkyl- C_{1-10} alkyl, cycloalkenyl, cycloalkenyl- C_{1-10} alkyl, aryl, aryl- C_{1-10} alkyl, heterocyclyl, heterocyclyl- C_{1-10} alkyl, heteroaryl, or heteroaryl- C_{1-10} alkyl; L^a is $-C(O)-$, $-O-C(O)-$, $-C(O)-O-$, $-C(O)-NR^c$, $-NR^c-C(O)-$, $-NR^c-C(O)-NR^d$, $-NR^c-C(O)-O-$, $-O-C(O)-NR^c$, $-S(O)_m-$, $-SO_2-NR^c$, $-NR^c-SO_2-$, $-O-$, $-NR^c$, or a bond (m , R^c and R^d have been defined above); and Z^2 is cycloalkyl, cycloalkyl- C_{1-10} alkyl, cycloalkenyl, cycloalkenyl- C_{1-10} alkyl, aryl, aryl- C_{1-10} alkyl, heterocyclyl, heterocyclyl- C_{1-10} alkyl, heteroaryl, heteroaryl- C_{1-10} alkyl or a bond. In one embodiment, Z^1 is cycloalkyl, cycloalkyl- C_{1-10} alkyl, aryl, aryl- C_{1-10} alkyl, heterocyclyl, heterocyclyl- C_{1-10} alkyl, heteroaryl, or heteroaryl- C_{1-10} alkyl; L^a is $-O-C(O)-$, $-C(O)-O-$, $-C(O)-NR^c$, $-NR^c-C(O)-$, $-SO_2-$, $-SO_2-NR^c$, $-NR^c-SO_2-$, $-O-$, $-NR^c$, or a bond; and Z^2 is aryl, aryl- C_{1-10} alkyl, heterocyclyl, heterocyclyl- C_{1-10} alkyl, or a bond. In one embodiment, Z^1 is aryl, aryl- C_{1-5} alkyl, heterocyclyl, heterocyclyl- C_{1-5} alkyl, heteroaryl, or heteroaryl- C_{1-5} alkyl; L^a is $-O-C(O)-$, $-C(O)-O-$, $-C(O)-NR^c$, $-NR^c-C(O)-$, $-SO_2-$, or a bond; and Z^2 is heterocyclyl, heterocyclyl- C_{1-5} alkyl, or a bond. In one embodiment, Z^1 is phenyl optionally substituted with Cy, $-CO-R^d$, halogen, oxo, aryl-substituted alkenyl; L^a is $-O-C(O)-$, $-C(O)-O-$, $-C(O)-NR^c$, $-NR^c-C(O)-$, or $-SO_2-$; and Z^2 is heterocyclyl or a bond.


In one embodiment, the compounds of this invention contain R^1 of formula (ii):



(ii)

wherein R^9 is C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, Cy, Cy- C_{1-10} alkyl, Cy- C_{2-10} alkenyl, or Cy- C_{2-10} alkynyl; each of R^{10} and R^{11} , independently, is hydrogen, aryl, alkyl, alkenyl or alkynyl, cycloalkyl, cycloalkenyl, or aryl-substituted alkyl; and R^{12} is H, C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, aryl, aryl- C_{1-10} alkyl, heteroaryl, or heteroaryl- C_{1-10} alkyl. Cy has the same definition as stated above. Each of alkyl, alkenyl and alkynyl is optionally substituted with one to four substituents independently selected from R^a , and aryl and heteroaryl are optionally substituted with one to four substituents independently selected from R^b . R^a and R^b have been defined above. Note that R^{11} , R^{12} and the carbon to which they are attached optionally form a 3-7 membered mono- or bicyclic ring containing 0-2 heteroatoms selected from N, O, and S.


In one embodiment, the compounds of this invention contain R^1 of formula (iii):




(iii)

wherein R^{14} is C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, Cy, Cy- C_{1-10} alkyl, Cy- C_{2-10} alkenyl, or Cy- C_{2-10} alkynyl; R^{15} is H, C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, aryl, aryl- C_{1-10} alkyl, heteroaryl, or heteroaryl- C_{1-10} alkyl; each of R^{16} , R^{17} , and R^{18} , independently, is H, C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, Cy, Cy- C_{1-10} alkyl, Cy- C_{2-10} alkenyl, Cy- C_{2-10} alkynyl, or a group selected from R^a . Cy has the same meaning as stated above (i.e., Cy represents cycloalkyl, heterocyclyl, aryl, or heteroaryl) is optionally substituted with one to four substituents independently selected from R^b or one of the following groups: $-NR^cC(O)NR^dSO_2R^d$, $-NR^cS(O)_mR^d$, $-OS(O)_2OR^c$, or $-OP(O)(OR^c)_2$. R^b has been defined above. Two of R^{16} , R^{17} , and R^{18} , when attached to a common ring atom, together with the

common ring atom optionally form a 5-7 membered saturated or unsaturated monocyclic ring containing zero to three heteroatoms selected from N, O, or S. Two of R¹⁶, R¹⁷, and R¹⁸, when attached to two adjacent ring atoms, together with these two ring atoms optionally form a 5-7 membered saturated or unsaturated monocyclic ring containing zero to three

- 5 heteroatoms selected from N, O, or S. The ring  represents a 3-7 membered saturated or unsaturated heterocyclyl or heteroaryl wherein each of Z, A, B₁ and B₂, independently, is a bond, -C-, -C-C-, -C=C-, a heteroatom selected from the group consisting of N, O, and S, or -S(O)_m- (with m being 0, 1, or 2). Y⁷ is -C(O)-, -C(O)O-, -C(O)NR^c-, -S(O)₂-, -P(O)(OR^c)-, or -C(O)-C(O)-. R^c has the same meaning as stated above. Each of the
- 10 alkyl, alkenyl and alkynyl is optionally substituted with one to four substituents independently selected from R^a, and each Cy is optionally substituted with one to four substituents independently selected from R^b. R^a and R^b have been defined above. In one


embodiment, the ring  in formula (ii), *supra*, represents azetidine, pyrrole, pyrrolidine, imidazole, pyrazole, triazole, pyridine, piperidine, pyrazine, piperazine,

15 pyrimidine, oxazole, thiazole, or morpholine. In one embodiment, the just-mentioned ring represents azetidine, pyrrole, pyrrolidine, imidazole, piperidine, or morpholine. In one embodiment, the just-mentioned ring represents pyrrolidine. In one embodiment, R¹⁵ is H or C₁₋₅ alkyl. In one embodiment, each of R¹⁶, R¹⁷, and R¹⁸, independently, is H, C₁₋₁₀ alkyl, Cy, -OR^c, -halogen, -S(O)_mR^c, -NR^cR^d, -NR^cC(O)R^d, -NR^cC(O)OR^d, -NR^cC(O)NR^dR^e, or oxo

20 (each of R^c, R^d, R^e, and m have been defined above). In one embodiment, Y⁷ is -O-C(O)-, -C(O)-O-, or -SO₂- (e.g., Y⁷ is -SO₂-). In one embodiment, R¹⁴ is Cy or Cy-C₁₋₅ alkyl (e.g., R¹⁴ is phenyl).

In one embodiment, the compounds of this invention contain L' having 2-4 (e.g., 2 or 3) carbon chain atoms.

- 25 In one embodiment, L' is of formula (iv):




(iv)

- wherein Y^1 is $-C(O)-$, $-O-C(O)-$, $-C(O)-O-$, $-C(O)-NR^c-$, $-NR^c-C(O)-$, $-NR^c-C(O)-NR^d-$, $-NR^c-C(O)-O-$, $-O-C(O)-NR^c-$, $-S(O)_m-$, $-S(O)_2-NR^c-$, $-NR^c-S(O)_2-$, $-NR^c-C(NR^m)-$, $-O-$, or $-NR^c-$ (R^c , R^d , R^m , and m have been defined above); R^2 is H, C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, Cy, Cy- C_{1-10} alkyl, Cy- C_{1-10} alkenyl, or Cy- C_{1-10} alkynyl; Y^2 is a bond or $-C(R^h)(R^i)-$, wherein each of R^h and R^i , independently, is H, C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, aryl, aryl- C_{1-10} alkyl, heteroaryl, or heteroaryl- C_{1-10} alkyl, and R^h and R^i can be taken together with the carbon to which they are attached to form a 3-7 membered ring containing 0-2 heteroatoms selected from N, O and S; X is $-C(O)OR^c$, $-P(O)(OR^c)(OR^d)$, $-P(O)(R^c)(OR^d)$, $-S(O)_mOR^c$, $-C(O)NR^cR^j$, or -5-tetrazolyl. m have been defined above. Each of said alkyl, alkenyl and alkynyl is optionally substituted with one to four substituents independently selected from R^a ; each of aryl and heteroaryl is optionally substituted with one to four substituents independently selected from R^b ; and Cy is a cycloalkyl, heterocyclyl, aryl, or heteroaryl. R^a and R^b have been defined above. Note that when Y^2 is not a bond, X is -
- COOH, $-COO-C_{1-4}$ alkyl, $-P(O)(OH)_2$, $-P(O)(OH)(O-C_{1-4}$ alkyl), $-P(O)(C_{1-4}$ alkyl) $_2$, $-P(O)(OH)(C_{1-4}$ alkyl), $-P(O)(O-C_{1-4}$ alkyl)(C_{1-4} alkyl), $-SO_2-C_{1-4}$ alkyl, $-CO-NH_2$, $-CO-NH(C_{1-4}$ alkyl), $-CO-N(C_{1-4}$ alkyl) $_2$, or -5-tetrazolyl. In one embodiment, Y^1 is $-NR^c-C(O)-$, $-NR^c-$, $-NR^c-S(O)_2-$, or $-NR^c-C(NR^m)-$. In one embodiment, Y^1 is $-NR^c-C(O)-$ (e.g., $-NH-CO-$ or $-N(C_{1-4}$ alkyl)- $CO-$; with the carbonyl group attaching to R^1). In one embodiment, R^2 is H or C_{1-5} alkyl. In one embodiment, R^2 is H. In one embodiment, Y^2 is a bond or $-C(R^h)(R^i)-$, wherein each of R^h and R^i , independently, is H or C_{1-5} alkyl. In one embodiment, Y^2 is a bond or $-CH_2-$. In one embodiment, X is $-C(O)OR^c$ (e.g., $-COOH$ or $-COO-C_{1-5}$ alkyl such as $-COO-CH_3$ or $-COO-CH_2CH_3$) or $-C(O)NR^cR^j$. In one embodiment, Y^1 is $-NR^c-C(O)-$ (e.g., $-NH-CO-$); R^2 is H or C_{1-5} alkyl (e.g., H); Y^2 is a bond or $-CH_2-$ (e.g., a bond); and X is -
- $C(O)OR^c$ where each R^c is independently H or C_{1-5} alkyl.


In one embodiment, the compounds of this invention contain L having 4-10 (e.g., 4-8 or 4-6) carbon chain atoms.

In one embodiment, L is of formula (v):



wherein Y^3 is a bond, C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, aryl, aryl- C_{1-10} alkyl, heteroaryl, or heteroaryl- C_{1-10} alkyl; and Y^4 is a bond, -C(O)-, -O-C(O)-, -C(O)-O-, -C(O)-NR^c-, -NR^c-C(O)-, -NR^c-C(O)-NR^d-, -NR^c-C(O)-O-, -O-C(O)-NR^c-, -S(O)_m-, -S(O)₂-NR^c-, -NR^c-S(O)₂-, -NR^c-C(NR^d)-, -O-, or -NR^c- (R^c, R^d, and m have been defined above). Each of alkyl, alkenyl, and alkynyl is optionally containing (interrupted by or terminally attached to) one to four heteroatoms selected from N, O, S, and -S(O)_m; and each of alkyl, alkenyl and alkynyl is optionally substituted with one to four substituents independently selected from R^a. Each of aryl and heteroaryl is optionally substituted with one to four substituents independently selected from R^b. R^a, R^b, R^c, R^d, and m have been defined above. Note that each of Y^3 and Y^4 is not a bond simultaneously. In one embodiment, Y^3 is a bond, C_{1-5} alkyl, or C_{1-5} alkenyl (e.g., Y^3 is a bond or C_{1-5} alkyl); and Y^4 is a bond, -C(O)-NR^c-, -C(O)-, -NR^c-, or -O-, where R^c is H or C_{1-5} alkyl (e.g., Y^4 is -C(O)-NH-).

In one embodiment, the compounds of this invention contain R³ with the formula: Z³-L^b-Z⁴-, wherein Z³ is Cy, Cy- C_{1-10} alkyl, Cy- C_{1-10} alkenyl, or Cy- C_{1-10} alkynyl; L^b is -C(O)-, -O-C(O)-, -C(O)-O-, -C(O)-NR^c-, -NR^c-C(O)-, -NR^c-C(O)-NR^d-, -NR^c-C(O)-O-, -O-C(O)-NR^c-, -S(O)_m-, -SO₂-NR^c-, -NR^c-SO₂-, -O-, -NR^c-, or a bond (R^c, R^d, and m have been defined above); and Z⁴ is cycloalkyl, cycloalkyl- C_{1-10} alkyl, cycloalkenyl, cycloalkenyl- C_{1-10} alkyl, aryl, aryl- C_{1-10} alkyl, heterocyclyl, heterocyclyl- C_{1-10} alkyl, heteroaryl, heteroaryl- C_{1-10} alkyl or a bond; or R³ is a moiety of formula (i):



each of m, R^c, R^d, R⁴, R⁵, R⁶, and Y⁵ have been defined in claim 1. In one embodiment, R⁴ is Z⁵-L^c-Z⁶-, wherein Z⁵ is Cy, Cy- C_{1-10} alkyl, Cy- C_{1-10} alkenyl, or Cy- C_{1-10} alkynyl; L^c is -C(O)-, -O-C(O)-, -C(O)-O-, -C(O)-NR^c-, -NR^c-C(O)-, -NR^c-C(O)-NR^d-, -NR^c-C(O)-O-, -O-C(O)-NR^c-, -S(O)_m-, -SO₂-NR^c-, -NR^c-SO₂-, -O-, -NR^c-, or a bond; and Z⁶ is cycloalkyl,

cycloalkyl-C₁₋₁₀ alkyl, cycloalkenyl, cycloalkenyl-C₁₋₁₀ alkyl, aryl, aryl-C₁₋₁₀ alkyl, heterocyclyl, heterocyclyl-C₁₋₁₀ alkyl, heteroaryl, heteroaryl-C₁₋₁₀ alkyl or a bond. R^c, R^d, m have been defined above. In one embodiment, each of Z³ and Z⁵, independently, is aryl, aryl-C₁₋₁₀ alkyl, aryl-C₁₋₁₀ alkenyl, aryl-C₁₋₁₀ alkynyl, heteroaryl, heteroaryl-C₁₋₁₀ alkyl, heteroaryl-C₁₋₁₀ alkenyl, or heteroaryl-C₁₋₁₀ alkynyl; each of L^b and L^c, independently, is -C(O)-, -S(O)_m-, -O-C(O)-, -C(O)-O-, -C(O)-NR^c-, -NR^c-C(O)-, -NR^c-C(O)-NR^d-, -SO₂-NR^c-, -NR^c-SO₂-, -O-, -NR^c-, or a bond; and each of Z⁴ and Z⁶, independently, is aryl, aryl-C₁₋₁₀ alkyl, heterocyclyl, heterocyclyl-C₁₋₁₀ alkyl, heteroaryl, heteroaryl-C₁₋₁₀ alkyl, or a bond. In one embodiment, each of Z³ and Z⁵, independently, is aryl, aryl-C₁₋₁₀ alkyl, heteroaryl, or heteroaryl-C₁₋₁₀ alkyl; each of L^b and L^c, independently, is -C(O)-, -SO₂-, -C(O)-NR^c-, -NR^c-C(O)-, or -NR^c-C(O)-NR^d-; where each of R^c and R^d, independently, is H or C₁₋₅ alkyl; and each of Z⁴ and Z⁶, independently, is aryl, aryl-C₁₋₁₀ alkyl, heterocyclyl, heterocyclyl-C₁₋₁₀ alkyl, heteroaryl, heteroaryl-C₁₋₁₀ alkyl, or a bond. In one embodiment, each of Z³ and Z⁵, independently, is aryl (e.g., phenyl or naphthyl); each of L^b and L^c, independently, is -NR^c-C(O)-NR^d- (e.g., -NH-CO-NH-, -N(methyl)-CO-NH-, or -NH-CO-N(methyl)-); and each of Z⁴ and Z⁶, independently, is aryl (e.g., phenyl or naphthyl). In one embodiment, Y⁵ is -CO- or -O-CO- (e.g., -CO-). In one embodiment, R⁵ is H or C₁₋₅ alkyl (e.g., H, methyl, or ethyl). In one embodiment, R⁶ is an amino acid side chain selected from the group consisting of cyclohexylalanine, leucine, isoleucine, allo-isoleucine, tert-leucine, norleucine, phenylalanine, phenylglycine, alanine, norvaline, valine, and 2-aminobutyric acid. In one embodiment, R⁶ is an amino acid side chain selected from the group consisting of leucine, isoleucine, allo-isoleucine, tert-leucine, norleucine, alanine, norvaline, valine, and 2-aminobutyric acid. In one embodiment, R⁶ is the side chain of leucine or isoleucine.


In one embodiment, the compounds of formula (I) contain R¹ with the formula Z¹-L^a-Z²-, wherein Z¹ is aryl (e.g., phenyl) optionally substituted with Cy, -CO-R^d, halogen, oxo, or aryl-substituted alkenyl; L^a is -O-C(O)-, -C(O)-O-, -C(O)-NR^c-, -NR^c-C(O)-, or -SO₂- (e.g., -SO₂-); and Z² is a bond, heteroaryl, heterocyclyl (e.g., azetidine, pyrrole, pyrrolidine, imidazole, piperidine, or morpholine); L' with formula (iv), *supra*, wherein Y¹ is -NR^c-C(O)-, -NR^c-, -NR^c-S(O)₂-, or -NR^c-C(NR^d)₂-; R² is H or C₁₋₅ alkyl; Y² is a bond or -C(R^h)(Rⁱ)₂-; and X is -C(O)OR^c; where each of R^c, R^h, and Rⁱ, independently, is H or C₁₋₅

alkyl (e.g., Y^1 is $-NH-C(O)-$; R^2 is H; Y^2 is a bond; and X is $-C(O)OH$); L with formula (v),
supra, wherein Y^3 is a bond, C_{1-5} alkyl, or C_{1-5} alkenyl; and Y^4 is a bond, $-C(O)-NR^c$,
 $-C(O)-$, $-NR^c$, or $-O-$, where R^c is H or C_{1-5} alkyl (e.g., Y^3 is a bond or C_{1-5} alkyl and Y^4 is
 $-C(O)-NH-$); and R^3 with the formula $Z^3-L^b-Z^4$ or formula (i), *supra*. When R^3 is of formula
 5 (i), R^4 is $Z^5-L^c-Z^6$, wherein Z^5 is aryl, aryl- C_{1-10} alkyl, aryl- C_{1-10} alkenyl, aryl- C_{1-10} alkynyl,
 heteroaryl, heteroaryl- C_{1-10} alkyl, heteroaryl- C_{1-10} alkenyl, or heteroaryl- C_{1-10} alkynyl; L^c is
 $-C(O)-$, $-S(O)_m-$, $-O-C(O)-$, $-C(O)-O-$, $-C(O)-NR^c$, $-NR^c-C(O)-$, $-NR^c-C(O)-NR^d$, $-SO_2-NR^c$,
 $-NR^c-SO_2-$, $-O-$, $-NR^c$, or a bond, with R^c and R^d , independently, being H or C_{1-5} alkyl; and
 Z^6 is aryl, aryl- C_{1-10} alkyl, heterocyclyl, heterocyclyl- C_{1-10} alkyl, heteroaryl, heteroaryl- C_{1-10}
 10 alkyl, or a bond. In one embodiment, Z^5 is aryl (e.g., phenyl or naphthyl); L^c is $-NR^c-C(O)-$
 $-NR^d$ (e.g., $-NH-CO-NH-$ or $-NH-CO-N(methyl)-$); and Z^6 is aryl (e.g., phenyl or naphthyl).
 In one embodiment, R^4 is o-methylphenyl-ureido-phenyl- CH_2- . In one embodiment, Y^5 is
 $-CO-$ or $-O-CO-$ (e.g., $-CO-$). In one embodiment, R^5 is H or C_{1-2} alkyl. In one embodiment,
 R^6 is an amino acid side chain selected from the group consisting of leucine, isoleucine, allo-
 15 isoleucine, tert-leucine, norleucine, alanine, norvaline, valine, and 2-aminobutyric acid (e.g.,
 leucine or isoleucine).

In one embodiment, the compounds of formula (I) contain R^1 with formula (ii), *supra*,
 wherein R^9 is C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, Cy, Cy- C_{1-10} alkyl, Cy- C_{2-10} alkenyl, or
 Cy- C_{2-10} alkynyl (e.g., aryl or heteroaryl); each of R^{10} and R^{11} , independently, is hydrogen,
 20 aryl, alkyl, alkenyl or alkynyl, cycloalkyl, cycloalkenyl, or aryl-substituted alkyl (e.g., H,
 alkyl, cycloalkyl, heterocyclyl, aryl, or heteroaryl); and R^{12} is H, C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10}
 alkynyl, aryl, aryl- C_{1-10} alkyl, heteroaryl, or heteroaryl- C_{1-10} alkyl (e.g., H, alkyl, alkenyl,
 alkynyl, heterocyclyl, or aryl). Cy has the same definition as stated above. Each of alkyl,
 alkenyl and alkynyl is optionally substituted with one to four substituents independently
 25 selected from R^a , and aryl and heteroaryl are optionally substituted with one to four
 substituents independently selected from R^b (e.g., halogen). R^a and R^b have been defined
 above. Note that R^{11} , R^{12} and the carbon to which they are attached optionally form a 3-7
 membered mono- or bicyclic ring containing 0-2 heteroatoms selected from N, O, and S. In
 this embodiment, the compounds also contain L' with formula (iv), *supra*, wherein Y^1 is
 30 $-NR^c-C(O)-$, $-NR^c$, $-NR^c-S(O)_2-$, or $-NR^c-C(NR^d)-$; R^2 is H or C_{1-5} alkyl; Y^2 is a bond or
 $-C(R^h)(R^i)-$; and X is $-C(O)OR^c$; where each of R^c , R^h , and R^i , independently, is H or C_{1-5}

alkyl (e.g., Y^1 is $-NH-C(O)-$; R^2 is H; Y^2 is a bond; and X is $-C(O)OH$); and L with formula (v), *supra*, wherein Y^3 is a bond, C_{1-5} alkyl, or C_{1-5} alkenyl; and Y^4 is a bond, $-C(O)-NR^c$, $-C(O)-$, $-NR^c$, or $-O-$, where R^c is H or C_{1-5} alkyl (e.g., Y^3 is a bond or C_{1-5} alkyl and Y^4 is $-C(O)-NH-$); and R^3 with the formula $Z^3-L^b-Z^4$ or formula (i), *supra*. When R^3 is of formula (i), R^4 is $Z^5-L^c-Z^6$, wherein Z^5 is aryl, aryl- C_{1-10} alkyl, aryl- C_{1-10} alkenyl, aryl- C_{1-10} alkynyl, heteroaryl, heteroaryl- C_{1-10} alkyl, heteroaryl- C_{1-10} alkenyl, or heteroaryl- C_{1-10} alkynyl; L^c is $-C(O)-$, $-S(O)_m-$, $-O-C(O)-$, $-C(O)-O-$, $-C(O)-NR^c$, $-NR^c-C(O)-$, $-NR^c-C(O)-NR^d$, $-SO_2-NR^c$, $-NR^c-SO_2-$, $-O-$, $-NR^c$, or a bond, with R^c and R^d , independently, being H or C_{1-5} alkyl; and Z^6 is aryl, aryl- C_{1-10} alkyl, heterocyclyl, heterocyclyl- C_{1-10} alkyl, heteroaryl, heteroaryl- C_{1-10} alkyl, or a bond. In one embodiment, Z^5 is aryl (e.g., phenyl or naphthyl); L^c is $-NR^c-C(O)-NR^d$ (e.g., $-NH-CO-NH-$ or $-NH-CO-N(methyl)-$); and Z^6 is aryl (e.g., phenyl or naphthyl). In one embodiment, R^4 is o-methylphenyl-ureido-phenyl- CH_2- . In one embodiment, Y^5 is $-CO-$ or $-O-CO-$ (e.g., $-CO-$). In one embodiment, R^5 is H or C_{1-2} alkyl. In one embodiment, R^6 is an amino acid side chain selected from the group consisting of leucine, isoleucine, allo-
 15 isoleucine, tert-leucine, norleucine, alanine, norvaline, valine, and 2-aminobutyric acid (e.g., leucine or isoleucine).

In one embodiment, the compounds of formula (I) contain R^1 with formula (iii), *supra*, wherein R^{14} is Cy or Cy- C_{1-5} alkyl (e.g., R^{14} is phenyl); R^{15} is H or C_{1-5} alkyl; each of R^{16} , R^{17} , and R^{18} , independently, is H, C_{1-10} alkyl, Cy, $-OR^c$, -halogen, $-S(O)_mR^c$, $-NR^cR^d$, $-NR^cC(O)R^d$, $-NR^cC(O)OR^d$, $-NR^cC(O)NR^dR^e$, or oxo (two of R^{16} , R^{17} , and R^{18} , when
 20 attached to two adjacent ring atoms, together with these two ring atoms optionally form a 5-7

membered cycloalkyl, heterocyclyl, aryl or heteroaryl); the ring  represents
 azetidine, pyrrole, pyrrolidine, imidazole, piperidine, or morpholine (e.g., pyrrolidine); Y^7 is $-O-C(O)-$, $-C(O)-O-$, or $-SO_2-$ (e.g., Y^7 is $-SO_2-$). The compounds also contain L' with
 25 formula (iv), *supra*, wherein Y^1 is $-NR^c-C(O)-$, $-NR^c$, $-NR^c-S(O)_2-$, or $-NR^c-C(NR^d)-$; R^2 is H or C_{1-5} alkyl; Y^2 is a bond or $-C(R^h)(R^i)-$; and X is $-C(O)OR^c$; where each of R^c , R^h , and R^i , independently, is H or C_{1-5} alkyl (e.g., Y^1 is $-NH-C(O)-$; R^2 is H; Y^2 is a bond; and X is $-C(O)OH$); and L with formula (v), *supra*, wherein Y^3 is a bond, C_{1-5} alkyl, or C_{1-5} alkenyl; and Y^4 is a bond, $-C(O)-NR^c$, $-C(O)-$, $-NR^c$, or $-O-$, where R^c is H or C_{1-5} alkyl (e.g., Y^3 is a

bond or C₁₋₅ alkyl and Y⁴ is -C(O)-NH-; and R³ with the formula Z³-L^b-Z⁴- or formula (i), *supra*. When R³ is of formula (i), R⁴ is Z⁵-L^c-Z⁶-, wherein Z⁵ is aryl, aryl-C₁₋₁₀ alkyl, aryl-C₁₋₁₀ alkenyl, aryl-C₁₋₁₀ alkynyl, heteroaryl, heteroaryl-C₁₋₁₀ alkyl, heteroaryl-C₁₋₁₀ alkenyl, or heteroaryl-C₁₋₁₀ alkynyl; L^c is -C(O)-, -S(O)_m-, -O-C(O)-, -C(O)-O-, -C(O)-NR^c-, -NR^c-C(O)-, -NR^c-C(O)-NR^d-, -SO₂-NR^c-, -NR^c-SO₂-, -O-, -NR^c-, or a bond, with R^c and R^d, independently, being H or C₁₋₅ alkyl; and Z⁶ is aryl, aryl-C₁₋₁₀ alkyl, heterocyclyl, heterocyclyl-C₁₋₁₀ alkyl, heteroaryl, heteroaryl-C₁₋₁₀ alkyl, or a bond. In one embodiment, Z⁵ is aryl (e.g., phenyl or naphthyl); L^c is -NR^c-C(O)-NR^d- (e.g., -NH-CO-NH- or -NH-CO-N(methyl)-); and Z⁶ is aryl (e.g., phenyl or naphthyl). In one embodiment, R⁴ is o-methylphenyl-ureido-phenyl-CH₂-. In one embodiment, Y⁵ is -CO- or -O-CO- (e.g., -CO-). In one embodiment, R⁵ is H or C₁₋₂ alkyl. In one embodiment, R⁶ is an amino acid side chain selected from the group consisting of leucine, isoleucine, allo-isoleucine, tert-leucine, norleucine, alanine, norvaline, valine, and 2-aminobutyric acid (e.g., leucine or isoleucine).


In one embodiment, the compounds of the invention are of formula (I) wherein R¹ is aryl or heterocyclyl-SO₂-aryl (e.g., pyrrolidine-SO₂-phenyl optionally substituted with alkyl or halo such as chloro, bromo, or iodo); L' is of formula (iv), *supra*, wherein Y¹ is -NH-CO-, -NH-, or -NH-C(NR^m)-NH-, R² is H, Y² is a bond or -CH₂-, and X is COOH; L is of formula (v), *supra*, wherein Y³ is -(CH₂)₀₋₅-, and Y⁴ is -CO-NH-; and R³ is o-methylphenyl-ureido-phenyl-CH₂- or of formula (i), *supra*, wherein R⁴ is o-methylphenyl-ureido-phenyl-CH₂-, Y⁵ is -CO- or -O-CO- (e.g., -CO-), R⁵ is H or methyl, and R⁶ is the side chain of leucine or isoleucine.

In one embodiment, the compounds of the invention contain L' and L as linker moiety, preferably composed of a chain containing C, O, S, or N atoms which link R¹ and R³ and allow both R¹ and R³ to interact, preferably bind, the VLA-4 molecule.

In one embodiment, the compounds of the invention have two terminally-located moieties of the formula Z^a-L^a-Z^b-. Each of Z^a and Z^b, independently, is an optionally substituted Cy, and L^a is a bond, or a linker moiety connecting Z^a and Z^b and can contain -C(O)-, -O-C(O)-, -C(O)-O-, -C(O)-NR^c-, -NR^c-C(O)-, -NR^c-C(O)-NR^d-, -NR^c-C(O)-O-, -O-C(O)-NR^c-, -S(O)_m-, -S(O)₂-NR^c-, -NR^c-S(O)₂-, -NR^c-C(NR^d)-, -O-, or -NR^c-. By "terminally-located" is meant that the moiety is monovalently attached to the rest of the molecule.


In one embodiment, the compounds of the invention have an IC_{50} of 5 nM or below, 2 nM or below, 1 nM or below, or 0.5 nM or below. IC_{50} values can be determined by binding assays as described below or other known conventional methods. In one embodiment, the compounds of the invention have a % bound to the Mn activated form of VLA-4 molecules of 50% or higher, 75% or higher, 90% or higher, or 95% or higher. In one embodiment, the compounds of the invention have a % bound to the Ca/Mg activated form of VLA-4 molecules of 50% or higher, 75% or higher, 90% or higher, or 95% or higher. % bound to the VLA-4 molecules can be determined by biological assays as described below.

In one embodiment, the compounds of this invention are of formula (II):




wherein each of R^1 , Y^1 , Y^2 , X , Y^3 , Y^4 , and R^3 have been defined above.

In one embodiment, the compounds of this invention is of formula (III):




Each of R^{21} and R^{22} , independently, is Cy, $-OR^c$, $-NO_2$, -halogen, $-S(O)_mR^c$, $-SR^c$, $-S(O)_2OR^c$, $-S(O)_2NR^cR^d$, $-NR^cR^d$, $-O(CR^cR^f)_nNR^cR^d$, $-C(O)R^c$, $-CO_2R^c$, $-CO_2(CR^cR^f)_nCONR^cR^d$, $-OC(O)R^c$, $-CN$, $-C(O)NR^cR^d$, $-NR^cC(O)R^d$, $-OC(O)NR^cR^d$, $-NR^cC(O)OR^d$, $-R^cC(O)NR^dR^e$, $-CR^c(NOR^d)$, $-CF_3$, $-OCF_3$, oxo, C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, aryl- C_{1-10} alkyl, or heteroaryl- C_{1-10} alkyl; wherein each of alkyl, alkenyl, alkynyl, aryl, heteroaryl assignable to R^{21} or R^{22} is optionally substituted with a group independently selected from R^b . R^{23} is H, C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, aryl, aryl- C_{1-10} alkyl, heteroaryl, or heteroaryl- C_{1-10} alkyl; wherein each of alkyl, alkenyl and alkynyl assignable to R^{23} is optionally substituted with one to four substituents independently selected from R^a , and aryl and heteroaryl are optionally substituted with one to four substituents independently selected from R^b . R^a , R^b and R^e have been defined above.

In one embodiment, the compounds of this invention are of formula (IV):



wherein each of R^1 , Y^1 , R^2 , Y^2 , X , Y^3 , Y^4 , and R^3 have been defined above.


In one embodiment, the compounds of this invention are of formula (V):



5

wherein each of R^1 , Y^1 , Y^2 , X , Y^3 , Y^4 , R^6 , R^5 , Y^5 and R^4 have been defined above.


In one embodiment, the compounds of this invention are of formula (VI):



wherein each of R^1 , X , Y^3 , Y^4 , R^6 , R^5 , and R^4 have been defined above.

10


In one embodiment, the compounds of this invention are of formula (VII):



wherein each of R^1 , X , Y^3 , R^6 , R^5 , and R^4 have been defined above.

Set forth below are some examples of a compound of this invention. For convenience, the nitrogen atom and the carbon atom in the column " $N(R^5)-CH(R^6)$ "



represents the α -nitrogen and the α -carbon atoms of the amino acid as indicated. For example, an entry "Leu" indicates that R^5 is H and R^6 is isobutyl.
























CPDB	R2/R4	Y3	N(R3)CH8	Y4	Y3	Y2	Y1	R2	R1	X	C ₂₀ H ₁₁
5192	oMePUPCH2	-C(=O)-	N-Me-Leu	-C(=O)NH-	-CH2CH2-	-	-NH-C(=O)-	H		CO2H	S
5241	Bu	-OC(=O)-	N-Me-Leu	-C(=O)NH-	-CH2CH2-	-	-NH-C(=O)-	H		CO2H	S
5247		-C(=O)-	N-Me-Leu	-C(=O)NH-	-CH2CH2-	-	-NH-C(=O)-	H		CO2H	S
5252	CH3	-	-	-C(=O)NH-	-CH2CH2-	-	-NH-C(=O)-	H		CO2H	S
5253	oMePUPCH2	-C(=O)-	Leu	-C(=O)NH-	-CH2-	-	-NH-C(=O)-	H		CO2H	S
5286	CH3	-	-	-C(=O)NH-	-CH2-	-	-NH-C(=O)-	H		CO2H	S
5292	oMePUPCH2	-C(=O)-	N-Me-Leu	-C(=O)NH-	-CH2CH2-	-	-NH-C(=O)-	H		CO2H	S
5310	Bn	-	-	-OC(=O)NH-	-CH2-	-	-NH-C(=O)-	H		CO2H	S
5337	Bn	-	-	-OC(=O)NH-	-CH2-	-	-NH-C(=O)-	H		CO2H	S


























CPDs	R2/R4	Y3	NPSCHER	Y4	Y3	Y2	Y1	R2	R1	X	Comp ID
538		-CO-	Leu	-C(O)NH-	-CH2-	-	-NHCO-	H	CH3	CO2H	S
5420	CH3	-CO-	N-Me-Leu	-C(O)NH-	-CH2-	-	-NHCO-	H		CO2H	S
5430	Bn	-CO-	N-Me-Leu	-C(O)NH-	-CH2-	-	-NHCO-	H		CO2H	S
5450		-CO-	Leu	-C(O)NH-	-CH2-	-	-NHCO-	H		CO2H	S
5451		-CO-	Leu	-C(O)NH-	-CH2-	-	-NHCO-	H		CO2H	S
5743	2-Cl-Bn	-	-	-C(O)NH-	-CH2-	-	-NHCO-	H		CO2H	S
5750	Bn	-CO-	N-Me-Leu	-C(O)NH-	-CH2-	-	-NHCO-	H		CO2H	S
5751		-CO-	N-Me-Leu	-C(O)NH-	-CH2-	-	-NHCO-	H		CO2H	S
5752	CH3	-	-	-C(O)NH-		-	-NHCO-	H		CO2H	S














CPDs	R3R4	Y3	NR5/CHR6	Y4	Y3	Y2	Y1	R2	R1	X	Con R
9785	CH3	-	-	-CO2NH-	-CH2CH2-	-	-NHCO2-	H		CO2H	S
9800	CH3	-CO2-	Leu	-CO2NH-		-	-NHCO2-	H		CO2H	S
9801		-	-	-CO2NH-	-CH2CH2-	-	-NHCO2-	H		CO2H	S
9803	oMePUPCH2	-	-	-CO2NH-	-CH2CH2-	-	-NHCO2-	H		CO2H	S
9835		-	-	-CO2NH-	-CH2CH2-	-	-NHCO2-	H	CH3	CO2H	S
9858		-CO2-	N-Me-Leu	-CO2NH-	-CH2-	-	-NHCO2-	H		CO2H	S
9869		-CO2-	Phe	-CO2NH-	-CH2-	-	-NHCO2-	H		CO2H	S
9870		-CO2-	Met22	-CO2NH-	-CH2-	-	-NHCO2-	H		CO2H	S
9871		-CO2-	Leu	-CO2NH-	-CH2-	-	-NHCO2-	H		CO2H	S





CPD#	R3R4	Y5	M/R5CHR8	Y4	Y3	Y2	Y1	R2	R1	X	Cen In.
6606		-C(=O)-	Pro	-C(=O)NH-	-CH2CH2-	-	-NH-C(=O)-	H		CO2H	S
6627		-C(=O)-	Pro	-C(=O)NH-	-CH2-	-	-NH-C(=O)-	H		CO2H	S
6714		-C(=O)-	N-Me-Leu	-C(=O)NH-	-CH2-	-	-NH-C(=O)-	H		CO2H	S
6715		-C(=O)-	Pro	-C(=O)NH-	-CH2-	-	-NH-C(=O)-	H		CO2H	S
6716		-C(=O)-	Leu	-C(=O)NH-	-CH2-	-	-NH-C(=O)-	H		CO2H	S
7080		-C(=O)-	N-Me-Leu	-C(=O)NH-	-CH2CH2-	-	-NH-C(=O)-	H		CO2H	S
7081		-	-	-C(=O)NH-	-CH2CH2-	-	-NH-C(=O)-	H		CO2H	S
7083		-	-	-CO-		-	-NH-C(=O)-	H		CO2H	S
7082		-C(=O)-	N-Me-Leu	-C(=O)NH-	-CH2CH2-	-	-NH-C(=O)-	H		CO2H	S





CPD#	R3/R4	Y5	NIP5/CHR8	Y4	Y3	Y2	Y1	R2	R1	X	Com in
7103			N-Me-Leu	-CO(NH)-	-CH2-	-	-NHCO-	H		CO2H	S
7105			Dansyl-Lys	-CO(NH)-	-CH2-	-	-NHCO-	H		CO2H	S
7109			N-Me-Leu	-CO(NH)-	-CH2-	-	-NHCO-	H		CO2H	S
7111			-	-CO(NH)-	-CH2-	-	-NHCO-	H		CO2H	S
7115			N-Me-Leu	-CO(NH)-	-CH2-	-	-NHCO-	H		CO2H	S
7117			-	-CO(NH)-	-CH2-	-	-NHCO-	H	CH3	CO2H	S
7119			-	-CO(NH)-	-CH2-	-	-NHCO-	H	CH3	CO2H	S
7147			-	-CO(NH)-	-CH2-	-	-NHCO-	H		CO2H	S
7148			-	-CO(NH)-	-CH2-	-	-NHCO-	H		CO2H	S

CPD#	R3R4	Y3	NIR5CHR8	Y4	Y3	Y2	Y1	R2	R1	X	Con Is
7150	2-Cl-8n	-	-	-O-C(=O)NH-	-CH2H-	-	-NHClO-	Bn		CO2H	S
7155	6-Me-PU-CH2	-	-	-O-C(=O)NH-	-CH2H-	-	-NHClO-	H		CO2H	S
7156	6-Me-PU-CH2	-	-	-O-C(=O)NH-	-CH2H-	-	-NHClO-	Bn		CO2H	R/S
7157		-	-	-O-C(=O)NH-	-CH2H-	-	-NHClO-	Bn		CO2H	R/S
7158	CH3	-	-	-O-C(=O)NH-	-CH2H-	-	-NHClO-	Bn		CO2H	R/S
7168	Bn	-	N-Me-Leu	-O-C(=O)NH-	-CH2H-	-	-NHClO-	H		CO2H	S
7171		-O-C(=O)-	N-Me-Leu	-O-C(=O)NH-	-CH2-	-	-NHClO-	H		CO2H	S
7172		-O-C(=O)-	Pro	-O-C(=O)NH-	-CH2-	-	-NHClO-	H		CO2H	S
7175		-O-C(=O)-	MeO2	-O-C(=O)NH-	-CH2-	-	-NHClO-	H		CO2H	S





CPD#	R3/R4	Y5	H35CHR8	Y4	Y3	Y2	Y1	R2	R1	X	Con No.
7117		-C(=O)-	Leu	-C(=O)NH-	-CH2-	-	-NHClO-	H		CO2H	S
7181	OMePUPCH2	-C(=O)-	N-Me-Leu	-C(=O)NH-	-[CH2]2-	-	-NHClO-	H		CO2CH3	S
7200	OMePUPCH2	-C(=O)-	N-Me-Leu	-C(=O)NH-	-[CH2]2-	-	-NHClO-	H		CO2H	S
7231	H	-	-	-NH-	-[CH2]4-	-	-NHClO-	H		CO2H	S
7233		-	-	-NH-	-[CH2]4-	-	-NHClO-	H		CO2H	S
7234	OMePUPCH2	-C(=O)-	Leu	-NH-	-[CH2]4-	-	-NHClO-	H		CO2H	S
7235		-	-	-C(=O)NH-	-[CH2]4-	-	-NHClO-	H		CO2H	S
7236		-	-	-C(=O)NH-	-[CH2]4-	-	-NHClO-	H		CO2H	S
7241	OMePUPCH2	-	-	-C(=O)NH-	-[CH2]4-	-	-NHClO-	H		CO2H	S





















CPD#	R3R4	Y3	NIR3CHIR3	Y4	Y3	Y2	Y1	R2	R1	X	Con lg
7255	Bn	_C(O)-	Pro	_C(O)NH-	_CH2H-	-	_NH2(O)-	H		CO2H	S
7256	oMePUPCH2	_C(O)-	Pro	_C(O)NH-	_CH2H-	-	_NH2(O)-	H		CO2H	S
7257	Bn	_C(O)-	Leu	_C(O)NH-	_CH2H-	-	_NH2(O)-	H		CO2H	S
7258	oMePUPCH2	_C(O)-	N-Me-Leu	_C(O)NH-	_CH2H-	-	_NH2(O)-	H		CO2H	S
7259	Bn	_C(O)-	Leu	_C(O)NH-	_CH2H-	-	_NH2(O)-	H		CO2H	S
7260	oMePUPCH2	_C(O)-	N-Me-Leu	_C(O)NH-	_CH2H-	-	_NH2(O)-	H		CONHCH3	S
7261	oMePUPCH2	_C(O)-	Gly	_C(O)NH-	_CH2H-	-	_NH2(O)-	H		CO2H	S
7262		_C(O)-	Leu	_C(O)NH-	_CH2-	-	_NH2(O)-	H		CO2H	S
7263		_C(O)-	Pro	_C(O)NH-	_CH2-	-	_NH2(O)-	H		CO2H	S










CPD#	R3/R4	Y3	NR3/CH3	Y4	Y2	Y1	R2	R1	X	Con Jg
7316		-CO-	Leu	-CO-NH-	-CH2-	-NHCO-	H		CO2H	S
7317		-CO-	Pro	-CO-NH-	-CH2-	-NHCO-	H		CO2H	S
7328		-CO-	Leu	-CO-NH-	-[CH2]2-	-NHCO-	H		CO2H	S
7330		-CO-	Pro	-CO-NH-	-[CH2]2-	-NHCO-	H		CO2H	S
7332		-CO-	α-N-Me-Cβ-Lys	-CO-NH-	-[CH2]2-	-NHCO-	H		CO2H	S
7378		-CO-	N-Me-Gly	-CO-NH-	-[CH2]2-	-NHCO-	H		CO2H	S
7652		-CO-	N-Me-Leu	-CO-NH-	-[CH2]2-	-NHCO-	H		CO2H	S
7788		-	-	-CO-NH-		-NHCO-	H		CO2H	S
7789		-CO-	N-Me-Leu	-CO-NH-	-[CH2]2-	-NHCO-	H		CH2OH	S





CPD#	R3/R4	Y8	NIP5/CH8	Y4	Y3	Y2	Y1	R2	R1	X	Con No.
7553	6MeFUPCH2	-	-	-C(O)NH-	-(CH2)4-	CH2	-NHClO-	H		CO2H	S
7556		-	-	-C(O)NH-	-(CH2)4-	CH2	-NHClO-	H		CO2H	S
7557		-	-	-C(O)NH-	-(CH2)4-	CH2	-NHClO-	H		CO2H	S
8066	CH3	-	-	-C(O)NH-	-(CH2)4-	CH2	-NHClO-	H		CO2H	S
8067	Bn	-	-	-C(O)NH-	-(CH2)4-	CH2	-NHClO-	H		CO2H	S
8122	6MeFUPCH2	-	-	-C(O)NH-	-(CH2)4-	CH2	-NHClO-	H		CO2H	S
8123		-	-	-C(O)NH-	-(CH2)4-	CH2	-NHClO-	H		CO2H	S
8147		-	-	-C(O)NH-	-(CH2)4-	CH2	-NHClO-	H		CO2H	S
8203	6MeFUPCH2	-	-	-C(O)NH-	-(CH2)5-	-	-NHClO-	H		CO2H	S



CPD#	R3/R4	Y3	NH5/CHR8	Y4	Y3	Y2	Y1	R2	R1	X	Con Gr.
8208		-	-	-CONH-	-CH2H-	CH2	-NHClO-	H	CH3	CO2H	S
8209		-	-	-CONH-	-CH2H-	CH2	-NHClO-	H		CO2H	S
8210		-	-	-CONH-	-CH2H-	CH2	-NHClO-	H	CH3	CO2H	S
8211		-	-	-CONH-	-CH2H-	CH2	-NHClO-	H		CO2H	S
8212		-	-	-CONH-	-CH2H-	CH2	-NHClO-	H		CO2H	S
8221		-CO-	N-Me-Leu	-CONH-	-CH2H-	-	-NHClO-	H		CO2H	S
8230		-CO-	N-Me-Leu	-CONH-	-CH2H-	-	-NHClO-	H		CO2H	S
8231		-CO-	N-Me-Leu	-CONH-	-CH2H-	-	-NHClO-	H		CO2H	S
8234		-CO-	N-Me-Leu	-CONH-	-CH2H-	-	-NHClO-	H		CO2H	S


CPD#	R3/R4	Y5	N(R5)CHR8	Y4	Y3	Y2	Y1	R2	R1	X	Can No.
8395	oMePUPCH2	-C(O)-	N-Me-Leu	-C(O)NH-	-CH2CH2-	-	-NHCl(O)-	H		CO2H	S
8396	oMePUPCH2	-	-	-C(O)NH-		-	-NHCl(O)-	H		CO2H	S
8398	oMePUPCH2	-C(O)-	N-Me-Leu	-C(O)NH-	-CH2CH2-	-	-NHCl(O)-	H		CO2H	S
8399	oMePUPCH2	-C(O)-	N-Me-Leu	-C(O)NH-	-CH2CH2-	-	-NHCl(O)-	H		CO2H	S
8401	oMePUPCH2	-C(O)-	N-Me-Leu	-C(O)NH-	-CH2CH2-	-	-NHCl(O)-	H		CO2H	S
8402	oMePUPCH2	-C(O)-	N-Me-Leu	-C(O)NH-	-CH2CH2-	-	-NHCl(O)-	H		CO2H	R
8403	oMePUPCH2	-C(O)-	N-Me-Leu	-C(O)NH-	-CH2CH2-	-	-NHCl(O)-	H		CO2H	R
8405	oMePUPCH2	-	-	-C(O)NH-	-CH2CH2-	-	-NH-	H	H	CO2H	S
8408	oMePUPCH2	-C(O)-	D-Me-Leu	-C(O)NH-	-CH2CH2-	-	-NHCl(O)-	H		CO2H	R







CPO#	RUR#	Y6	NPR9CHR8	Y4	Y3	Y2	Y1	B2	R1	X	COSY JR.
8346	OMEPUFCH2	-C(=O)-	D-N-Me-Leu	-C(O)NH-	-[CH2]2-	-	-NH-C(=O)-	H		CO2H	R
8352	OMEPUFCH2	-	-	-C(O)NH-	-[CH2]4-	-	-NH-C(=O)-	H	tBu	CO2H	S
8354	4-OMEPUP	-	-	-O-	-[CH2]2-	-	-NH-C(=O)-	H		CO2H	R
8355	OMEPUFCH2	-	-	-C(O)NH-		-	-NH-C(=O)-	H		CO2H	R
8357	OMEPUFCH2	-C(=O)-	D-N-Me-Leu	-C(O)NH-	-[CH2]2-	-	-NH-C(=O)-	H		CO2H	S
8358	OMEPUFCH2	-C(=O)-	D-N-Me-Leu	-C(O)NH-	-[CH2]2-	-	-NH-C(=O)-	H		CO2H	S
8359		-	-	-C(O)NH-	-[CH2]2-	-	-NH-C(=O)-	H		CO2H	S
8446	OMEPUFCH2	-	-	-C(O)NH-	-[CH2]4-	-	-NH-C(=O)-	H	Bn	CO2H	S
8449	OMEPUFCH2	-	-	-C(O)NH-	-[CH2]4-	CH2	-NH-C(=O)-	H		CO2H	S


CPD#	R3/R4	Y5	N(R5)CHR8	Y4	Y3	Y2	Y1	R2	R1	X	Comp. ID.
B450	Bn	-	-	-CO ₂ NH-	-CH ₂ N-	-CH ₂ -	-NHClO-	H		CO ₂ H	S
B451		-CO ₂ -	Pro	-CO ₂ NH-	-CH ₂ N-	-CH ₂ -	-NHClO-	H		CO ₂ H	S
B452		-	-	-CO ₂ NH-	-CH ₂ N-	-CH ₂ -	-NHClO-	H		CO ₂ H	S
B453	oMePUPCH ₂	-	-	-CO ₂ NH-	-CH ₂ N-	-CH ₂ -	-NHClO-	H		CO ₂ H	S
B454		-CO ₂ -	Pro	-CO ₂ NH-	-CH ₂ N-	-CH ₂ -	-NHClO-	H		CO ₂ H	S
B455		-	-	-CO ₂ NH-	-CH ₂ N-	-CH ₂ -	-NHClO-	H		CO ₂ H	S
B456		-	-	-CO ₂ NH-	-CH ₂ N-	-CH ₂ -	-NHClO-	H		CO ₂ H	S
B457		-CO ₂ -	Pro	-CO ₂ NH-	-CH ₂ N-	-CH ₂ -	-NHClO-	H		CO ₂ H	S
B458	oMePUPCH ₂	-	-	-CO ₂ NH-	-CH ₂ N-	-CH ₂ -	-NHClO-	H		CO ₂ H	S
B459	Bn	-	-	-CO ₂ NH-	-CH ₂ N-	-CH ₂ -	-NHClO-	H		CO ₂ H	S





CPD#	R1R2R4	Y5	NH2CHR8	Y4	Y3	Y2	Y1	R2	R1	X	Con No.
8450		-C(=O)-	Pro	-C(=O)NH-	-(CH2)4-	CH2	-NH-C(=O)-	H		CO2H	5
8451		-	-	-C(=O)NH-	-(CH2)4-	CH2	-NH-C(=O)-	H		CO2H	5
8452	OMePUPCH2	-	-	-C(=O)NH-	-(CH2)4-	CH2	-NH-C(=O)-	H	OMePUPALeu	CO2H	5
8453	Bn	-	-	-C(=O)NH-	-(CH2)4-	CH2	-NH-C(=O)-	H	OMePUPALeu	CO2H	5
8454		-C(=O)-	Pro	-C(=O)NH-	-(CH2)4-	CH2	-NH-C(=O)-	H	OMePUPALeu	CO2H	5
8455		-C(=O)-	Pro	-C(=O)NH-	-(CH2)4-	CH2	-NH-C(=O)-	H	CH3	CO2H	5
8456		-	-	-C(=O)NH-	-(CH2)4-	CH2	-NH-C(=O)-	H	CH3	CO2H	5
8457	OMePUPCH2	-C(=O)-	Leu	-C(=O)NH-	-(CH2)2-	-	-NH-C(=O)-	H		CO2H	5
8458	2-OMePUP	-	-	-O-	-(CH2)2-	-	-NH-C(=O)-	H		CO2H	5



















CPDs	REMR	Y8	N(R)C(R)R8	Y4	Y3	Y2	Y1	R2	R1	X	Con DL
8485		-	-	-CO ₂ NH-		-	-NHClO-	H		CO ₂ H	S
8491		-CO ₂ -	Leu	-CO ₂ NH-	-CH ₂ -	-	-NHClO-	H		CO ₂ H	S
8493		-CO ₂ -	N-Me-Leu	-CO ₂ NH-	-CH ₂ -	-	-NHClO-	H		CO ₂ H	S
8494		-CO ₂ -	D-N-Me-Leu	-CO ₂ NH-	-CH ₂ -	-	-NHClO-	H		CO ₂ H	R
8513		-	-	-CO ₂ NH-	-CH ₂ -	CH ₂	-NHClO-	H		CO ₂ H	S
8514		-	-	-CO ₂ NH-		-	-NHClO-	H		CO ₂ H	S
8515		-	-	-CO ₂ NH-		-	-NHClO-	H		CO ₂ H	S
8516		-	-	-CO ₂ NH-		-	-NHClO-	H		CO ₂ H	S
8519		-	-	-CO ₂ NH-	-CH ₂ -	CH ₂	-NHClO-	H		CO ₂ H	S
















CPDs	R3/R4	Y5	NR3CHR8	Y4	Y3	Y2	Y1	R2	R1	X	Con SL
8520		-	-	-CO2NH-		-	-NHCO2-	H		CO2H	S
8528	αMePUPCH2	-CO2-	N-Me-Leu	-CO2NH-	-CH2-	-	-NHCO2-	H		CO2H	S
8532		-CO2-	Leu	-CO2NH-	-CH2-	-	-NHCO2-	H		CO2H	S
8533		-CO2-	Leu	-CO2NH-	-CH2-	-	-NHCO2-	H		CO2H	S
8534		-CO2-	Leu	-CO2NH-	-CH2-	-	-NHCO2-	H		CO2H	S
8535	αMePUPCH2	-CO2-	N-Me-Leu	-CO2NH-	-CH2-	-	-NHCO2-	H		CO2H	S
8537	αMePUPCH2	-	-	-CO2NH-		-	-NHCO2-	H		CO2H	S
8538	αMePUPCH2	-CO2-	Leu	-CO2NH-	-CH2-	-	-NHCO2-	H		CO2H	S
8539	αMePUPCH2	-CO2-	Leu	-CO2NH-	-CH2-	-	-NHCO2-	H		CO2H	S

































CPD#	R3R4	Y5	NR3CORR5	Y4	Y3	Y2	Y1	R2	R1	X	Com In
8566		-C(O)-	Leu	-C(O)NH-	-CH2CH2-	-	-NH-C(O)-	H		CO2H	S
8567		-C(O)-	Leu	-C(O)NH-	-CH2CH2-	-	-NH-C(O)-	H		CO2H	S
8571	OMePUPCH2	-C(O)-	N-Me-Leu	-C(O)NH-	-CH2CH2-	-	-NH-C(O)-	H		CO2H	S
8582	OMePUPCH2	-C(O)-	N-Me-Leu	-C(O)NH-	-CH2CH2-	-	-NH-C(O)-	H		CO2H	S
8583	OMePUPCH2	-C(O)-	N-Me-Leu	-C(O)NH-	-CH2CH2-	-	-NH-C(O)-	H		CO2H	S
8585		-C(O)-	Leu	-C(O)NH-	-CH2CH2-	-	-NH-C(O)-	H		CO2H	S
8586	OMePUPCH2	-C(O)-	N-Me-Leu	-C(O)NH-	-CH2CH2-	-	-NH-C(O)-	H		CO2H	S
8588	Bn	-	-	-C(O)NH-		-	-NH-C(O)-	H		CO2H	S
8587	OMePUPCH2	-	-	-C(O)NH-		-	-NH-C(O)-	H		CO2H	S



CPD#	R3/R4	V5	N/R3CH8	Y4	Y3	Y2	Y1	R2	R1	X	Comp ID
8320		-	-	-CO ₂ NH-	-CH ₂ -	-	-NHClO-	H		CO ₂ H	R
8321	OMePUPCH ₂	-CO-	Pro	-CO ₂ NH-	-CH ₂ -	-	-NHClO-	H		CO ₂ H	R
8325	OMePUPCH ₂	-CO-	N-Me-Leu	-CO ₂ NH-	-CH ₂ -	-	-NHClO-	H		CO ₂ H	S
8329	OMePUPCH ₂	-	-	-CO ₂ NH-	-CH ₂ -	-	-NHClO-	H		CO ₂ H	R
8330	OMePUPCH ₂	-CO-	Pro	-CO ₂ NH-	-CH ₂ -	-	-NHClO-	H		CO ₂ H	R
8332	Bz	-CO-	Pro	-CO ₂ NH-	-CH ₂ -	-	-NHClO-	H		CO ₂ H	R
8337		-	-	-CO ₂ NH-	-CH ₂ -	-	-NHClO-	H		CO ₂ H	R
8338	Bn	-CO-	Leu	-CO ₂ NH-	-CH ₂ -	-	-NHClO-	H		CO ₂ H	R
8339	Bn	-CO-	Leu	-CO ₂ NH-	-CH ₂ -	-	-NHClO-	H		CO ₂ H	R





CPD#	R3R4	Y5	MRSCH8	Y4	Y3	Y2	Y1	R2	R1	X	Com. #
8642			N-Me-Leu	-C(O)NH-	-CH2CH2-	-	-NHClO-	H		CO2H	S
8643			Asp	-C(O)NH-	-CH2CH2-	-	-NHClO-	H		CO2H	S
8646			-	-C(O)NH-		-CH2-	-NHClO-	H		CO2H	S
8656			-	-C(O)NH-	-CH2CH2-	-	-NHClO-	H		CO2H	R
8674			N-Me-Leu	-C(O)NH-	-CH2CH2-	-	-NHClO-	H		CO2H	S
8684			-	-C(O)NH-		-	-NHClO-	H		CO2H	S
8685			N-Me-Leu	-C(O)NH-	-CH2CH2-	-	-NHClO-	H		CO2H	S
8699			N-Me-Leu	-C(O)NH-	-CH2CH2-	-	-NHClO-	H		CO2H	S
8690			N-Me-Leu	-C(O)NH-	-CH2CH2-	-	-NHClO-	H		CO2H	S





















CPD#	R3R4	Y5	NR9CHR8	Y4	Y3	Y2	Y1	R2	R1	X	Gen ID
8698	oMePUPCH2	_COO-	N-Me-Glu	_C(O)NH-	_CH2CH2-	-	_NHCOO-	H		CO2H	S
8773	oMePUPCH2	_COO-	N-Me-Leu	_C(O)NH-	_CH2CH2-	-	_NHCOO-	H		CO2H	S
8746	oMePUPCH2	_COO-	N-Me-Leu	_C(O)NH-	_CH2CH2-	-	_NHCOO-	H		CO2H	S
8749	oMePUPCH2	_COO-	N-Me-Leu	_C(O)NH-	_CH2CH2-	-	_NHCOO-	H		CO2H	S
8758	oMePUPCH2	_COO-	N-Me-Leu	_C(O)NH-	_CH2CH2-	-	_NHCOO-	H		CO2H	S
8796	oMePUPCH2	_COO-	N-Me-Leu	_C(O)NH-	_CH2CH2-	-	_NHCOO-	H		CO2H	S
8797	oMePUPCH2	_COO-	N-Me-Leu	_C(O)NH-	_CH2CH2-	-	_NHCOO-	H		CO2H	S
8809	oMePUPCH2	_COO-	N-Me-Leu	_C(O)NH-	_CH2CH2-	-	_NHCOO-	H		CO2H	S
8805	oMePUPCH2	_COO-	Leu	_C(O)NH-	_CH2-	-	-	H		CO2H	R/S

CPD#	R3R4	Y5	N(R5)CHR8	Y4	Y3	Y2	Y1	R2	R1	X	Con Lg
8908	oMePUPCH2	_C(O)_	Pro	_C(O)NH-	_CH2-	-	-	H		CO2H	RS
8929	oMePUPCH2	_C(O)_	N-Me-Leu	_C(O)NH-	_CH2CH2-	-	_NHCO(O)-	H		CO2H	S
9120	oMePUPCH2	_C(O)_	N-Me-Leu	_C(O)NH-	_CH2CH2-	-	_NHCO(O)-	H		CO2H	S
9140	oMePUPCH2	_C(O)_	N-Me-Leu	_C(O)NH-	_CH2CH2-	-	_NHCO(O)-	H	_CH3	CO2H	S
9169	oMePUPCH2	_C(O)_	N-Me-Leu	_C(O)NH-	_CH2CH2-	-	_NHCO(O)-	H		CO2H	S
9170	oMePUPCH2	_C(O)_	N-Me-Leu	_C(O)NH-	_CH2CH2-	-	_NHCO(O)-	H		CO2H	S
9171	oMePUPCH2	_C(O)_	N-Me-Leu	_C(O)NH-	_CH2CH2-	-	_NHCO(O)-	H		CO2H	S
9182	oMePUPCH2	_C(O)_	N-Me-Leu	_C(O)NH-	_CH2CH2-	-	_NHCO(O)-	H		CO2H	S
9227	oMePUPCH2	_C(O)_	N-Me-Leu	_C(O)NH-	_CH2CH2-	-	_NHCO(O)-	H		CO2H	S









CPD#	BJRR	Y5	NIPASCHRB	Y4	Y3	Y2	Y1	R2	R1	X	Cont No.
9232	oAmPUPCH2	-	-	_C(O)NH-		-	_NHClO4	H		CO2H	S
9233	oAmPUPCH2	_ClO4	Leu	_C(O)NH-		-	_NHClO4	H		CO2H	S
9234	oAmPUPCH2	-	-	_C(O)NH-		-	_NHClO4	H		CO2H	S
9235	oAmPUPCH2	_ClO4	Leu	_C(O)NH-		-	_NHClO4	H		CO2H	S
9236	oAmPUPCH2	-	-	_C(O)NH-		-	_NHClO4	H		CO2H	S
9237	oAmPUPCH2	_ClO4	Leu	_C(O)NH-		-	_NHClO4	H		CO2H	S
9238	oAmPUPCH2	-	-	_C(O)NH-		-	_NHClO4	H		CO2H	S
9239	oAmPUPCH2	_ClO4	Leu	_C(O)NH-		-	_NHClO4	H		CO2H	S
9244	oAmPUPCH2	_ClO4	N-Me-Leu	_C(O)NH-	_CH2CH2-	-	_NHClO4	H		CO2H	S



CPD	R3R4	Y3	MP5CHR8	Y4	Y3	Y2	Y1	R2	R1	X	Com AL
9270	oMePUPCH2	-	-	-C(=O)-		-	-C(=O)-	H		CO2H	S
9271	oMePUPCH2	-C(=O)-	Leu	-C(=O)-		-	-C(=O)-	H		CO2H	S
9273	oMePUPCH2	-C(=O)-	Leu	-C(=O)-		-	-C(=O)-	H		CO2H	S
9274	oMePUPCH2	-	-	-C(=O)-		-	-C(=O)-	H		CO2H	S
9275	oMePUPCH2	-C(=O)-	Leu	-C(=O)-		-	-C(=O)-	H		CO2H	S
9276	oMePUPCH2	-	-	-C(=O)-		-	-C(=O)-	H		CO2H	S
9277	oMePUPCH2	-C(=O)-	Leu	-C(=O)-		-	-C(=O)-	H		CO2H	S
9215	oMePUPCH2	-C(=O)-	N-Me-Leu	-C(=O)NH-	-C(=O)2-	-	-NHC(=O)-	H		CO2H	S
9418	oMePUPCH2	-C(=O)-	N-Me-Leu	-C(=O)NH-	-C(=O)2-	-	-NHC(=O)-	H		CO2H	S



CPD#	R324	Y5	NIR5CHRB	Y4	Y3	Y2	Y1	R2	R1	X	Con lg.
9437	oMePUPCH2	-C(O)-	N-Me-Leu	-C(O)NH-	-(CH2)2-	-	-NHC(O)-	H		CO2H	S
9621		-C(O)-	Leu	-C(O)NH-	-(CH2)2-	-	-NHC(O)-	H		CO2H	S

Another aspect of this invention relates to the use of one or more of the inhibitors described above or a salt thereof for the manufacture of a medicament for treating the above-mentioned disorders.

A further aspect of this invention relates to a composition comprising a
5 pharmaceutical carrier and an effective amount of a compound of formula (I), *supra*.

Still a further aspect of this invention relates to a method of inhibiting VLA-4-dependent cell adhesion, comprising administering to a patient in need thereof an effective amount of a compound of formula (I), *supra*.

The ability of the compounds of this invention to antagonize the actions of VLA4
10 makes them useful for preventing, treating, or reversing the symptoms, disorders or diseases induced by the binding of VLA4 to its ligands. Thus these antagonists will inhibit cell adhesion processes including cell activation, migration, proliferation and differentiation. Accordingly, another aspect of the present invention provides methods for the treatment, prevention, alleviation, or suppression of diseases or disorders mediated by the VLA4
15 pathway. Such diseases and disorders include, for example, asthma, multiple sclerosis, allergic rhinitis, allergic conjunctivitis, inflammatory lung diseases, rheumatoid arthritis, septic arthritis, type I diabetes, organ transplant rejection, inflammatory bowel disease, and others.

Compounds of the invention contain one or more asymmetric centers and thus can
20 occur as racemates and racemic mixtures, single enantiomers, diastereomeric mixtures and individual diastereomers. The present invention is meant to comprehend all such isomeric forms of the compounds of the invention.

The claimed invention is also intended to encompass pharmaceutically acceptable salts of Formula I. The term "pharmaceutically acceptable salts" refers to salts prepared
25 from pharmaceutically acceptable non-toxic bases or acids including inorganic or organic bases and inorganic or organic acids. Salts derived from inorganic bases include aluminum, ammonium, calcium, copper, ferric, ferrous, lithium, magnesium, manganic salts, manganous, potassium, sodium, zinc, and the like. Particularly preferred are the ammonium, calcium, magnesium, potassium and sodium salts.

30 Salts derived from pharmaceutically acceptable organic non-toxic bases include salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring

substituted amines, cyclic amines, and basic ion exchange resins, such as arginine, betaine, caffeine, choline, N,N'-dibenzylethylenediamine, diethylamine, 2-diethylaminoethanol, 2-dimethylaminoethanol, ethanolamine, ethylenediamine, N-ethyl-morpholine, N-ethylpiperidine, glucamine, glucosamine, histidine, hydrabamine, isopropylamine, lysine, 5 methylglucamine, morpholine, piperazine, piperidine, polyamine resins, procaine, purines, theobromine, triethylamine, trimethylamine, tripropylamine, tromethamine, and the like.

When the compound of the present invention is basic, salts may be prepared from pharmaceutically acceptable non-toxic acids, including inorganic and organic acids. Such acids include acetic, benzenesulfonic, benzoic, camphorsulfonic, citric, ethanesulfonic, 10 fumaric, gluconic, glutamic, hydrobromic, hydrochloric, isethionic, lactic, maleic, malic, mandelic, methanesulfonic, mucic, nitric, pamoic, pantothenic, phosphoric, succinic, sulfuric, tartaric, p-toluenesulfonic acid, and the like. Particularly preferred are citric, hydrobromic, hydrochloric, maleic, phosphoric, sulfuric and tartaric acids.

As used herein, the term "alkyl," alone or in combination, refers to a straight-chain or 15 branched-chain alkyl radical containing from 1 to 10, preferably from 1 to 6 and more preferably from 1 to 4, carbon atoms. Examples of such radicals include, but are not limited to, methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, pentyl, isoamyl, hexyl, decyl and the like.

The term "alkenyl," alone or in combination, refers to a straight-chain or branched- 20 chain alkenyl radical containing from 2 to 10, preferably from 2 to 6 and more preferably from 2 to 4, carbon atoms. Examples of such radicals include, but are not limited to, ethenyl, E- and Z-propenyl, isopropenyl, E- and Z-butenyl, E- and Z-isobutenyl, E- and Z-pentenyl, decenyl and the like.

The term "alkynyl," alone or in combination, refers to a straight-chain or branched- 25 chain alkynyl radical containing from 2 to 10, preferably from 2 to 6 and more preferably from 2 to 4, carbon atoms. Examples of such radicals include, but are not limited to, ethynyl (acetylenyl), propynyl, propargyl, butynyl, hexynyl, decynyl and the like.

The term "hydrocarbon linker moiety" refers to an alkylene moiety which may contain one or more double or triple bonds. For example, L can be 3-methyloctylene (i.e., a 30 straight chain containing 8 carbon chain atoms) interrupted by, or terminally attached to, an amide linkage (-NH-CO-).

The term "cycloalkyl," alone or in combination, refers to a cyclic alkyl radical containing from 3 to 8, preferably from 3 to 6, carbon atoms. Examples of such cycloalkyl radicals include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and the like.

5 The term "cycloalkenyl," alone or in combination, refers to a cyclic carbocycle containing from 4 to 8, preferably 5 or 6, carbon atoms and one or more double bonds. Examples of such cycloalkenyl radicals include, but are not limited to, cyclopentenyl, cyclohexenyl, cyclopentadienyl and the like.

10 The term "aryl" refers to a carbocyclic aromatic group selected from the group consisting of phenyl, naphthyl, indenyl, indanyl, azulenyl, fluorenyl, and anthracenyl; or a heterocyclic aromatic group selected from the group consisting of furyl, thienyl, pyridyl, pyrrolyl, oxazolyly, thiazolyl, imidazolyl, pyrazolyl, 2-pyrazolynyl, pyrazolidinyl, isoxazolyl, isothiazolyl, 1,2,3-oxadiazolyl, 1,2,3-triazolyl, 1,3,4-thiadiazolyl, pyridazinyl, pyrimidinyl, pyrazinyl, 1,3,5-triazinyl, 1,3,5-trithianyl, indolizynyl, indolyl, isoindolyl, 3H-indolyl, 15 indolynyl, benzo[b]furanyl, 2,3-dihydrobenzofuranyl, benzo[b]thiophenyl, 1H-indazolyl, benzimidazolyl, benzthiazolyl, purinyl, 4H-quinolizynyl, quinolynyl, isoquinolynyl, cinnolynyl, phthalazinyl, quinazolinyl, quinoxalinyl, 1,8-naphthyridinyl, pteridinyl, carbazolyl, acridinyl, phenazinyl, phenothiazinyl, and phenoxazinyl.

20 "Aryl" groups, as defined in this application may independently contain one to three substituents which are independently selected from the group consisting of hydrogen, halogen, hydroxyl, amino, nitro, trifluoromethyl, trifluoromethoxy, alkyl, alkenyl, alkynyl, cyano, carboxy, carboalkoxy, Ar'-substituted alkyl, Ar'-substituted alkenyl or alkynyl, 1,2-dioxymethylene, 1,2-dioxyethylene, alkoxy, alkenoxy or alkynoxy, Ar'-substituted alkoxy, Ar'-substituted alkenoxy or alkynoxy, alkylamino, alkenylamino or alkynylamino, Ar'- 25 substituted alkylamino, Ar'-substituted alkenylamino or alkynylamino, Ar'-substituted carbonyloxy, alkylcarbonyloxy, aliphatic or aromatic acyl, Ar'-substituted acyl, Ar'-substituted alkylcarbonyloxy, Ar'-substituted carbonylamino, Ar'-substituted amino, Ar'-substituted oxy, Ar'-substituted carbonyl, alkylcarbonylamino, Ar'-substituted alkylcarbonylamino, alkoxy-carbonylamino, Ar'-substituted alkoxycarbonyl-amino, Ar'- 30 oxycarbonylamino, alkylsulfonylamino, mono- or bis-(Ar'-sulfonyl)amino, Ar'-substituted alkyl-sulfonylamino, morpholinocarbonylamino, thiomorpholinocarbonylamino, N-alkyl

guanidino, N-Ar' guanidino, N-N-(Ar',alkyl) guanidino, N,N-(Ar',Ar')guanidino, N,N-dialkyl
 guanidino, N,N,N-trialkyl guanidino, N-alkyl urea, N,N-dialkyl urea, N-Ar' urea, N,N-
 (Ar',alkyl) urea and N,N-(Ar')₂ urea; wherein "Ar'" is a carbocyclic or heterocyclic aryl
 group as defined above having one to three substituents selected from the group consisting of
 5 hydrogen, halogen, hydroxyl, amino, nitro, trifluoromethyl, trifluoromethoxy, alkyl, alkenyl,
 alkynyl, 1,2-dioxymethylene, 1,2-dioxyethylene, alkoxy, alkenoxy, alkynoxy, alkylamino,
 alkenylamino or alkynylamino, alkylcarbonyloxy, aliphatic or aromatic acyl,
 alkylcarbonylamino, alkoxycarbonylamino, alkylsulfonylamino, N-alkyl or N,N-dialkyl urea.

The term "alkoxy," alone or in combination, refers to an alkyl ether radical, wherein
 10 the term "alkyl" is as defined above. Examples of suitable alkyl ether radicals include, but
 are not limited to, methoxy, ethoxy, n-propoxy, iso-propoxy, n-butoxy, iso-butoxy, sec-
 butoxy, tert-butoxy and the like.

The term "alkenoxo," alone or in combination, refers to a radical of formula alkenyl-
 O-, wherein the term "alkenyl" is as defined above provided that the radical is not an enol
 15 ether. Examples of suitable alkenoxy radicals include, but are not limited to, allyloxy, E- and
 Z-3-methyl-2-propenoxy and the like. The term "alkynyloxy", alone or in combination, refers
 to a radical of formula alkynyl-O-, wherein the term "alkynyl" is as defined above provided
 that the radical is not an ynol ether. Examples of suitable alkynoxy radicals include, but are
 not limited to, propargyloxy, 2-butynyloxy and the like.

20 The term "thioalkoxy" refers to a thioether radical of formula alkyl-S-, wherein alkyl
 is as defined above.

The term "alkylamino," alone or in combination, refers to a mono- or di-alkyl-
 substituted amino radical (i.e., a radical of formula alkyl-NH- or (alkyl)₂-N-), wherein the
 term "alkyl" is as defined above. Examples of suitable alkylamino radicals include, but are
 25 not limited to, methylamino, ethylamino, propylamino, isopropylamino, t-butylamino, N,N-
 diethylamino and the like.

The term "alkenylamino," alone or in combination, refers to a radical of formula
 alkenyl-NH- or (alkenyl)₂N-, wherein the term "alkenyl" is as defined above, provided that
 the radical is not an enamine. An example of such alkenylamino radicals is the allylamino
 30 radical.

The term "alkynylamino," alone or in combination, refers to a radical of formula alkynyl-NH- or (alkynyl)₂N-, wherein the term "alkynyl" is as defined above, provided that the radical is not an ynamine. An example of such alkynylamino radicals is the propargyl amino radical.

5 The term "aryloxy," alone or in combination, refers to a radical of formula aryl-O-, wherein aryl is as defined above. Examples of aryloxy radicals include, but are not limited to, phenoxy, naphthoxy, pyridyloxy and the like.

 The term "arylamino," alone or in combination, refers to a radical of formula aryl-NH-, wherein aryl is as defined above. Examples of arylamino radicals include, but are not limited to, phenylamino (anilido), naphthylamino, 2-, 3- and 4-pyridylamino and the like.

10 The term "biaryl," alone or in combination, refers to a radical of formula aryl-aryl-, wherein the term "aryl" is as defined above.

 The term "thioaryl," alone or in combination, refers to a radical of formula aryl-S-, wherein the term "aryl" is as defined above. An example of a thioaryl radical is the thiophenyl radical.

15 The term "aryl-fused cycloalkyl," alone or in combination, refers to a cycloalkyl radical which shares two adjacent atoms with an aryl radical, wherein the terms "cycloalkyl" and "aryl" are as defined above. An example of an aryl-fused cycloalkyl radical is the benzo-fused cyclobutyl radical.

20 The term "aliphatic acyl," alone or in combination, refers to radicals of formula alkyl-CO-, alkenyl-CO- and alkynyl-CO-derived from an alkane-, alkene- or alkynycarboxylic acid, wherein the terms "alkyl", "alkenyl" and "alkynyl" are as defined above. Examples of such aliphatic acyl radicals include, but are not limited to, acetyl, propionyl, butyryl, valeryl, 4-methylvaleryl, acryloyl, crotyl, propiolyl, methylpropiolyl and the like.

25 The term "aromatic acyl," alone or in combination, refers to a radical of formula aryl-CO-, wherein the term "aryl" is as defined above. Examples of suitable aromatic acyl radicals include, but are not limited to, benzoyl, 4-halobenzoyl, 4-carboxybenzoyl, naphthoyl, pyridylcarbonyl and the like.

 The terms "morpholinocarbonyl" and "thiomorpholinocarbonyl," alone or in combination with other terms, refer to an N-carbonylated morpholino and an N-carbonylated thiomorpholino radical, respectively.

The term "alkylcarbonylamino," alone or in combination, refers to a radical of formula alkyl-CONH-, wherein the term "alkyl" is as defined above.

The term "alkoxycarbonylamino," alone or in combination, refers to a radical of formula alkyl-OCONH-, wherein the term "alkyl" is as defined above.

5 The term "alkylsulfonylamino," alone or in combination, refers to a radical of formula alkyl-SO₂NH-, wherein the term "alkyl" is as defined above.

The term "arylsulfonylamino," alone or in combination, refers to a radical of formula aryl-SO₂NH-, wherein the term "aryl" is as defined above.

10 The term "N-alkylurea," alone or in combination, refers to a radical of formula alkyl-NH-CO-NH-, wherein the term "alkyl" is as defined above.

The term "N-arylurea," alone or in combination, refers to a radical of formula aryl-NH-CO-NH-, wherein the term "aryl" is as defined above.

The term "halogen" means fluorine, chlorine, bromine and iodine.

15 The term "leaving group" generally refers to groups readily displaceable by a nucleophile, such as an amine, and alcohol or a thiol nucleophile. Such leaving groups are well known and include carboxylates, N-hydroxysuccinimide, N-hydroxybenzotriazole, halogen (halides), triflates, tosylates, mesylates, alkoxy, thioalkoxy and the like.

20 The terms "activated derivative of a suitably protected α -amino acid" and "activated substituted-phenylacetic acid derivative" refer to the corresponding acyl halides (e.g. acid fluoride, acid chloride and acid bromide), corresponding activated esters (e.g. nitrophenyl ester, the ester of 1-hydroxybenzotriazole, HOBT, or the ester of hydroxysuccinimide, HOSu), and other conventional derivatives within the skill of the art.

As used throughout this application, the term "patient" refers to mammals, including humans. And the term "cell" refers to mammalian cells, including human cells.

25 In view of the above definitions, other chemical terms used throughout this application can be easily understood by those of skill in the art. Terms may be used alone or in any combination thereof. The preferred and more preferred chain lengths of the radicals apply to all such combinations.

30 Other features or advantages of the present invention will be apparent from the following detailed description of several embodiments, and also from the appending claims.

DETAILED DESCRIPTION

Compounds of this invention may be synthesized using any conventional technique, several of which are exemplified herein. Preferably, these compounds are chemically synthesized from readily available starting materials, such as α -amino acids and their functional equivalents. Modular and convergent methods for the synthesis of these compounds are also preferred. In a convergent approach, for example, large sections of the final product are brought together in the last stages of the synthesis, rather than by incremental addition of small pieces to a growing molecular chain.

Compounds of the invention, $R^3-L-L'-R^1$, according to one embodiment, can be represented as $R^3-Y^4-Y^3-CH(X)-Y^1-R^1$. This compound can be viewed as a dipeptide derivative: with R^1 as an amino acid residue or a derivative thereof; Y^1 as an amide linkage, or a derivative thereof, between the two residues; X as a carboxylate or a derivative thereof; C as the α -carbon atom of the second residue; and $R^3-Y^4-Y^3-$ as the side chain of the second residue.

In the general method illustrated below, the compound $R^3-Y^4-Y^3-CH(X)-Y^1-R^1$ is prepared by first coupling a properly protected $Y^4'-Y^3-CH(X)-Y^1'$ with a properly protected R^3' . Y^3 and X have been defined above. Y^4' , Y^1' , and R^3' are precursors of Y^4 , Y^1 , and R^3 , respectively.

Compounds of this invention may be synthesized using any conventional technique, several of which are exemplified herein. Preferably, these compounds are chemically synthesized from readily available starting materials, such as α -amino acids and their functional equivalents. Modular and convergent methods for the synthesis of these compounds are also preferred. In a convergent approach, for example, large sections of the final product are brought together in the last stages of the synthesis, rather than by incremental addition of small pieces to a growing molecular chain.

Compounds of the invention, $R^3-L-L'-R^1$, according to one embodiment, can be represented as $R^3-Y^4-Y^3-CH(X)-Y^1-R^1$. This compound can be viewed as a dipeptide derivative: with R^1 as an amino acid residue or a derivative thereof; Y^1 as an amide linkage, or a derivative thereof, between the two residues; X as a carboxylate or a derivative thereof; C as the α -carbon atom of the second residue; and $R^3-Y^4-Y^3-$ as the side chain of the second residue.

In the general method illustrated below, the compound $R^3-Y^4-Y^3-CH(X)-Y^1-R^1$ is prepared by first coupling a properly protected $Y^4-Y^3-CH(X)-Y^1$ with a properly protected R^3 . Y^3 and X have been defined above. Y^4 , Y^1 , and R^3 are precursors of Y^4 , Y^1 , and R^3 , respectively.

- 5 Compounds of the formula $Y^4-Y^3-CH(X)-Y^1$ are available commercially or can be prepared according to methods known one of ordinary skill in the art. For example, if Y^1 is an amino group; X is a carboxylate; and Y^4-Y^3 is $NH_2-(CH_2)_3-$, the compound $Y^4-Y^3-CH(X)-Y^1$ is ornithine. As another example, if Y^1 is an amino group, X is carboxylate and Y^4-Y^3 is 4- NH_2 -phenyl- CH_2- , the compound $Y^4-Y^3-CH(X)-Y^1$ is 4-aminophenylalanine, available by reduction of commercially available 4-nitrophenylalanine. Further reduction of the phenyl moiety produces a compound wherein Y^1 is an amino group, X is carboxylate and Y^4-Y^3 is 4- NH_2 -cyclohexyl- CH_2- , or 4-aminocyclohexylalanine, available commercially as a mixture of *cis* and *trans* isomers. As mentioned above, proper protecting groups are required to prevent certain functionalities from undergoing undesired reactions.
- 10 Using ornithine as an example, Y^1 and X are functionalities that are not involved in the first coupling reaction, and should be protected with common amino protecting groups such as carbamates (e.g., *t*-butyl carbamate (BOC) and benzyl carbamate (CBZ)) and common carboxyl protecting groups such as substituted esters (e.g., ethyl ester and methoxymethyl ester). For more appropriate protecting groups, see T. W. Greene, Protecting Groups in Organic Synthesis, John Wiley & Sons, New York, 1981, and references cited therein.
- 15 The compound R^3 can be represented by the formula $Z^3-L^b-Z^4-T$ or $R^4-Y^5-N(R^5)-CH(R^6)-T'$. Each of T and T' is a functionality which joins with Y^4 to form Y^4 . For example, if the desired Y^4 is an amide linkage, it can be formed by reacting an amine group (Y^4) with a carboxyl group (T or T') in the presence of a common coupling reagent such as
- 20 benzotriazol-1-yloxytris(dimethylamino)-phosphonium hexafluorophosphate (BOP) or *O*-benzo-triazol-1-yl-*N,N,N',N'*-tetramethyluronium hexafluorophosphate (HBTU). As another example, if the desired Y^4 is an aryl ether, it can be formed by reacting a phenol with an alcohol in the presence of diethylazodicarboxylate (DEAD) and triphenylphosphine.


- When R^3 is of the formula $Z^3-L^b-Z^4-T$, the compound is available commercially or can be prepared according to methods known one of ordinary skill in the art. For example, when Z^3 is 2-methyl phenyl; Z^4 is phenylmethyl; L^b is $-NH-CO-NH-$ and T is $-COOH$, R^3 is
- 30

o-methylphenyl-ureido-phenyl acetic acid and can be obtained by reaction of 4-aminophenylacetic acid with 2-methylphenyl isocyanate. As another example, when Z^3 is 3-indole; Z^4 is phenylmethyl; L^b is $-\text{CO}-\text{NH}-$ and T is $-\text{COOH}$, $R^{3'}$ is 3-indolecarboxamido-phenyl acetic acid and can be obtained by reaction of 4-aminophenylacetic acid with indole-
 5 3-carbonyl chloride.

When $R^{3'}$ is of the formula $R^4-Y^5-N(R^5)-CH(R^6)-T'$, $Y^{4'}-Y^3-CH(X)-Y^{1'}$ can couple to $NH(R^5)-CH(R^6)-T'$ to form the intermediate $NH(R^5)-CH(R^6)-Y^4-Y^3-CH(X)-Y^{1'}$ prior to further coupling to $R^4-Y^{5'}$ to form $R^4-Y^5-N(R^5)-CH(R^6)-Y^4-Y^3-CH(X)-Y^{1'}$. $Y^{5'}$ is a functionality which, upon undergoing further coupling reactions, gives rise to the
 10 functionality Y^5 . Note that the compound $NH(R^5)-CH(R^6)-T'$ can be an amino acid derivative which is commercially available and can be prepared using conventional methods by one of ordinary skill in the art. For example, when T' is carboxyl; R^6 is isobutyl; and R^5 is methyl, the compound $NH(R^5)-CH(R^6)-T'$ is N-methylleucine. $R^4-Y^{5'}$ can be coupled to $NH(R^5)-CH(R^6)-Y^4-Y^3-CH(X)-Y^{1'}$ by commonly used synthetic methods. For example, if
 15 $Y^{5'}$ is carboxyl, the resulting Y^5 is an amide linkage and can be prepared using common peptide synthesis reagents as mentioned above. As another example, if $Y^{5'}$ is an halide or sulfonate the resulting Y^5 is a secondary or tertiary amine resulting from alkylation of the starting amine. Alternatively, to form the compound $R^4-Y^5-N(R^5)-CH(R^6)-Y^4-Y^3-CH(X)-Y^{1'}$, $NH(R^5)-CH(R^6)-T'$ can first couple to $R^4-Y^{5'}$ to form the intermediate $R^4-Y^5-N(R^5)-CH(R^6)-T'$ prior to further coupling to $Y^{4'}-Y^3-CH(X)-Y^{1'}$. Example 1 below provides a
 20 detailed procedure wherein $R^{3'}$ is of the formula $R^4-Y^5-N(R^5)-CH(R^6)-$.

Alternatively, when $R^{3'}$ is of the formula $Z^3-L^b-Z^4-T$, it can react with $Y^{4'}-Y^3-CH(X)-Y^{1'}$ to form $Z^3-L^b-Z^4-Y^4-Y^3-CH(X)-Y^{1'}$. See Example 2.


The final product $R^3-Y^4-Y^3-CH(X)-Y^1$ can then be formed by reacting either $R^4-Y^5-N(R^5)-CH(R^6)-Y^4-Y^3-CH(X)-Y^{1'}$ or $Z^3-L^b-Z^4-Y^4-Y^3-CH(X)-Y^{1'}$ with $R^{1'}$ (the precursor of R^1). The moiety Y^1 can be formed in a similar manner as Y^4 .
 25



A cell adhesion inhibitor of the invention can be purified by conventional methods such as chromatography or crystallization.

Set forth below are five general methods for preparing a compound of this invention.

5 General Method A – Solid-Phase Preparation of Diaminopropionate Derivatives:



Orthogonally Fmoc/Dde Protected Wang Resin (II): *S*-N- α -Fmoc-N- β -Dde-diaminopropionic acid, **I** (4.95 g, 10.1 mmol), was attached to Wang resin (7.88 g, 0.64 mmol/g, 100-200 mesh) by reaction with 2,6-dichlorobenzoyl chloride (1.45 mL, 10.1 mmol) and dry pyridine (1.35 mL) in 40 mL dry DMF. The mixture was shaken for 16 h at room temperature. The resin was isolated by filtration and was washed three times each with DMF and dichloromethane. The resin was capped by reaction with dichlorobenzoyl chloride and pyridine (2 mL each) for 2 h followed by washing as above. The resulting resin contained 0.64 mmol/g Fmoc as determined by piperidine treatment and measurement of A_{290} .


Deprotection and Acylation of N- α : The diaminopropionate resin, II, was treated with 20% piperidine in DMF for 15 min after which it was filtered and washed with DMF and dichloromethane. The deprotected resin was immediately acylated by treatment with R^1CO_2H (2 eq), HATU (2 eq) and diisopropylethylamine (4 eq). The reactions were shaken
5 for 2 h, filtered and the acylation was repeated. Completion of acylation was determined by a negative Kaiser test. The resin was filtered and washed with DMF and dichloromethane. If R^1CO_2H is an Fmoc protected amino acid, the deprotection and acylation are repeated as described above.

Deprotection and Acylation of N- β : The acylated diaminopropionate resin, III, was
10 treated with 2% hydrazine in DMF for 1 h, after which it was filtered and washed with DMF and dichloromethane. The deprotected resin was immediately acylated by treatment with R^3CO_2H (2 eq), HATU (2 eq) and diisopropylethylamine (4 eq). The reactions were shaken for 2 h, filtered and the acylation was repeated. The resin was filtered and washed with DMF and dichloromethane.

15 Cleavage of Final Product from Resin: The diacyl diaminopropionate resin, IV, was treated with 95% TFA/5% water for 1 h. The solvent was removed by filtration and the resin was washed with two small portions of TFA. The combined TFA solutions were concentrated under vacuum and the resulting residue was purified by reverse-phase hplc yielding pure diacyldiaminopropionate derivatives.

20

General Method B - Preparation of beta-Lysine Derivatives:



Omega-N-Cbz-beta-N-BOC-beta-homolysine Methyl Ester (II): Omega-N-Cbz-beta-N-BOC-beta-homolysine, I, was dissolved in N,N-dimethylformamide. To this solution was added sodium bicarbonate (10 equivalents) and then iodomethane (6 equivalents) with stirring. After stirring overnight at room temperature, the reaction mixture was partitioned between water and ethyl acetate. The organic layer was washed with saturated sodium chloride solution, then dried over sodium sulfate. Filtering and evaporation of the solvent was followed by silica gel chromatography (hexane/ethyl acetate) to yield ester II.

Beta-N-BOC-beta-homolysine Methyl Ester (III): N-Cbz carbamate II was dissolved in methanol. To this was added 10% palladium on carbon. The mixture was flushed with nitrogen, then hydrogen (50 psi) was added. After stirring overnight, the catalyst was removed using a Whatman PTFE filter and the solution was concentrated to yield crude amine III.


N-omega Acylation: Amine III (111 mg), 2-(1H-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (HBTU, 1.1 equivalents) and $\text{R}^1\text{CO}_2\text{H}$ (1.1 equivalents) were dissolved in N,N-dimethylformamide. To this solution was added N,N-

diisopropylethylamine (2.5 equivalents). After stirring overnight, the reaction was quenched with 5% aqueous citric acid solution, then extracted with ethyl acetate. The organics were washed with saturated sodium chloride solution, then dried over sodium sulfate. Filtration and removal of the solvent by rotary evaporation yielded crude amide IV, which was used without further purification.

N-beta Deprotection and Acylation: Crude N-BOC carbamate IV was treated with saturated hydrogen chloride in ethyl acetate, prepared by bubbling hydrogen chloride gas through cold (zero degree) ethyl acetate solution for 30 minutes. The reaction was stirred for one hour, then concentrated to dryness to yield crude amine V, which was used without further purification. Crude amine V was dissolved in N,N-dimethylformamide along with R^3CO_2H (1 equivalent) and HBTU (1.1 equivalent). With stirring was added N,N-diisopropylethylamine (7.5 equivalents). After stirring overnight, the reaction was partitioned between 5% aqueous citric acid and ethyl acetate. The organic layer was washed with saturated sodium chloride solution, then dried over sodium sulfate. Filtration of the drying agent and evaporation of the solvent gave crude amide VI, which was used without further purification.

Final Deprotection: Methyl ester VI was dissolved in 1:1 tetrahydrofuran and methanol. With stirring was added aqueous lithium hydroxide (2 N). After stirring for one hour, the reaction mixture was concentrated to dryness. The residue was partitioned between 1 N aqueous hydrogen chloride and ethyl acetate, and the organic layer was washed with saturated sodium chloride. Drying over sodium sulfate, filtering and evaporating gave crude acid. Purification by preparative reverse-phase high performance liquid chromatography gave pure acid.

General Method C – Solid-Phase Preparation of Lysine Derivatives:



- Fmoc/Dde Lysine Wang Resin (II): N-α-Fmoc-N-β-Dde-Lysine, I (5.0 g, 9.39 mmol), was attached to Wang resin (7.34 g, 0.64 mmol/g, 100-200 mesh) by reaction with 2,6-dichlorobenzoyl chloride (1.33 mL, 10.1 mmol) and dry pyridine (1.27 mL) in 50 mL dry
- 5 DMF. The mixture was shaken for 16 h at room temperature. The resin was isolated by filtration and was washed three times each with DMF and dichloromethane. The resin was capped by reaction with dichlorobenzoyl chloride and pyridine (2 mL each) for 2 h followed


by washing as above. The resulting resin contained 0.56 mmol/g Fmoc as determined by piperidine treatment and measurement of A_{290} .

Deprotection and Acylation of N- α : The diaminopropionate resin, II, was treated with 20% piperidine in DMF for 15 min after which it was filtered and washed with DMF and dichloromethane. The deprotected resin was immediately acylated by treatment with R^1CO_2H (2 eq), HATU (2 eq) and diisopropylethylamine (4 eq). The reactions were shaken for 2 h, filtered and the acylation was repeated. Completion of acylation was determined by a negative Kaiser test. The resin was filtered and washed with DMF and dichloromethane. If R^1CO_2H is an Fmoc protected amino acid, the deprotection and acylation are repeated as described above.

Deprotection and Acylation of N- ϵ : The acylated lysine resin, III, was treated with 2% hydrazine in DMF for 1 h, after which it was filtered and washed with DMF and dichloromethane. The deprotected resin was immediately acylated by treatment with R^3CO_2H (2 eq), HATU (2 eq) and diisopropylethylamine (4 eq). The reactions were shaken for 2 h, filtered and the acylation was repeated. The resin was filtered and washed with DMF and dichloromethane.

Cleavage of Final Product from Resin: The diacyl lysine resin, IV, was treated with 95% TFA/5% water for 1 h. The solvent was removed by filtration and the resin was washed with two small portions of TFA. The combined TFA solutions were concentrated under vacuum and the resulting residue was purified by reverse-phase HPLC yielding pure diacyllysine derivatives.

General Method D: Preparation of oMePUPA-N-MeLeu- α,γ -diaminobutyric Acid Derivatives:



N- α -CBZ-L-2,4-diaminobutyric acid methyl ester hydrochloride (I): In a 500 mL RB flask was suspended 8.4 g (33.3 mmol) N- α -CBZ-L-2,4-diaminobutyric acid in 200 mL methanol with stirring. This was cooled to 0°C (ice bath), and then 14.6 mL (200 mmol) SOCl₂ was added dropwise over 15 minutes to give a colorless solution. The solution was
5 allowed to warm to RT and stirred overnight. The solution was concentrated, redissolved in MeOH and concentrated 2x, then dissolved in CH₂Cl₂, concentrated, and placed under high vacuum for 16 hours to give compound I as a slightly yellow foam, massing to 10.33 g (34.2 mmol, 103%). M/z = 267.1 (M+H⁺).

BOC-N-methyl-Leuciny-(N- α -CBZ)-GABA methyl ester (II): In a 500mL RB
10 flask was dissolved 10.33 g (33.3 mmol) of I (MW = 302) in 100 mL dry dimethylformamide (DMF) with stirring to give a colorless solution. To this was added 17.4 mL (100 mmol) of diisopropylethylamine (DIEA), then 7.96 g (32.5 mmol) of Boc-N-Me-Leucine, and finally 14.83 g (39.0 mmol) of O-(7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (HATU) to give a yellow solution. This was stirred overnight, after
15 which HPLC showed no starting material. The solution was diluted with ethyl acetate (EtOAc, 500mL) and washed with 1N HCl (2x), 1N NaOH (2x), and brine (1x). The organic phase was dried over anhydrous MgSO₄, filtered, and concentrated to a red oil. Chromatography with 2:1 hexanes/EtOAc vs. silica gave 12.56 g (25.5 mmol, 78%) of II (R_f = 0.46 with 1:1 Hex/EtOAc vs. silica) as a yellow syrup (HPLC, >99%). M/z = 494.3
20 (M+H⁺).

H-N-methyl-Leuciny-(N- α -CBZ)-GABA methyl ester trifluoroacetate salt (III): In a 50 mL RB flask was dissolved 0.50 g (1.01 mmol) of II (MW=493) in 10 mL CH₂Cl₂ with stirring to give a colorless solution. To this was added 2 mL (26 mmol, large excess) of trifluoroacetic acid and the resulting solution was stirred for four hours, after which HPLC
25 showed no starting material. The solution was concentrated, redissolved in CH₂Cl₂ and concentrated (2x), then placed under high vacuum overnight to give 0.52 g (~ quantitative) of III as a very pale yellow oil. M/z = 394.4 (M+H⁺). Material carried through.

oMePUPA-N-methyl-Leuciny-(N- α -CBZ)-GABA methyl ester (IV): In a 10 mL vial was dissolved 0.52 g (1.01 mmol) of III (MW=507) in 5 mL DMF with stirring to give a
30 pale yellow solution. To this was added 525 μ L (3.0 mmol) of DIEA, then 284 mg (1.0 mmol) of oMePUPA free acid (Ricerca; MW=284), and finally 0.42 g (1.1 mmol) of HATU

to give a yellow solution. This was stirred overnight, after which HPLC showed no starting material remaining. The solution was diluted with EtOAc (75 mL) and washed with 1N HCl (3x), 1N NaOH (3x), and brine (1x). The organic phase was dried with MgSO₄, filtered, and the filtrate concentrated to a yellow oil/solid mixture. Chromatography with 1:2
5 acetonitrile/CH₂Cl₂ vs. silica gave 0.49 g (0.74 mmol, 74%) of VI (R_f = 0.56 with 1:1 acetonitrile/CH₂Cl₂ vs. silica) as a bright white, foamy solid (HPLC, >99%). M/z = 660.1 ($M+H^+$).


oMePUPA-N-methyl-Leuciny-(N- α -H)-GABA methyl ester Hydrochloride (V): In an 85 mL high-pressure vessel was dissolved 400 mg (0.61 mmol) of IV (MW=659) in 10
10 mL MeOH with stirring to give a colorless solution. The vessel was flushed with nitrogen, and ~50mg (catalytic) of 10% palladium on carbon was added. The sides of the vessel were washed with additional MeOH, and the vessel capped with a hydrogenation head. The vessel was charged with 60 psi H₂ and the mixture stirred overnight, after which the vessel was purged to ambient atmosphere. The mixture was filtered through Celite 545, the filter pad
15 washed with additional (10 mL) MeOH, and the filtrate concentrated. The residue was dissolved in minimal (2 mL) MeOH and dripped into ice-cold 1.0M HCl in diethyl ether to give a white precipitate. The solid was triturated in the HCl/ether for 20 minutes, then filtered, the solid washed with ether, and air-dried for one hour. The white solid was then crushed into a powder with a spatula, washed with additional ether, and air-dried overnight to
20 give 336 mg (0.60 mmol, 98%) of V as a white powder (HPLC, >99%). ESMS m/z = 526.6 ($M+H^+$).

Acylation and final hydrolysis: Crude amine V was dissolved in N,N-dimethylformamide along with R³CO₂H (1 equivalent) and HBTU (1.1 equivalent). With stirring was added N,N-diisopropylethylamine (4 equivalents). After stirring overnight, the
25 reaction was partitioned between 5% aqueous citric acid and ethyl acetate. The organic layer was washed with saturated sodium chloride solution, then dried over sodium sulfate. Filtration of the drying agent and evaporation of the solvent gave crude amide, which could be purified by reverse-phase hplc. Methyl ester was dissolved in 1:1 tetrahydrofuran and methanol. With stirring was added aqueous lithium hydroxide (2 N). After stirring for one
30 hour, the reaction mixture was concentrated to dryness. The residue was partitioned between 1 N aqueous hydrogen chloride and ethyl acetate, and the organic layer was washed with

saturated sodium chloride. Drying over sodium sulfate, filtering and evaporating gave crude acid. Purification by preparative reverse-phase high performance liquid chromatography gave pure product.

5

General Method E - Solution-Phase Synthesis from Diamino Acids:



The orthogonally N- α -Boc / Cbz protected diamine, **I**, was converted to methyl ester **II** by reaction with methyl iodide (5 eq) and potassium carbonate (5 eq) in acetone at room temperature for 16 h. The reaction mixture was diluted with water and extracted with ethyl acetate. The organics were washed with water, saturated sodium bicarbonate and brine, dried over sodium sulfate and filtered. Product was eluted through silica in ethyl acetate and hexanes.

N-alpha deprotection and acylation: The fully protected diamine, II, was dissolved in 3N HCl in EtOAc and was stirred 1 h at room temperature. The solution was concentrated under reduced pressure. The resulting solid was suspended in diethyl ether, isolated by filtration, washed with ether and dried under vacuum. The hydrochloride, III, thus isolated
5 was treated with HATU (1.25 eq), diisopropylethylamine (4 eq) and R^1CO_2H (1.25 eq) in dry DMF, and was stirred under nitrogen for 16 h. The reaction mixture was diluted with 5% citric acid and was extracted with EtOAc. . The organics were washed with water, saturated sodium bicarbonate and brine, dried over sodium sulfate and filtered. The solution was concentrated under reduced pressure and the residue was purified by elution through silica in
10 EtOAc and hexane, providing pure product, IV.

Distal nitrogen deprotection and acylation: The CBz protected intermediate, IV, was dissolved in methanol and was degassed. 10% Pd on activated carbon was added and the mixture was stirred under 60 psi hydrogen for 3 to 16 h. The reaction was filtered and concentrated. The resulting free amine was immediately acylated by reacting with HATU
15 (1.25 eq), diisopropylethylamine (4 eq) and R^3CO_2H (1.25 eq) in dry DMF, with stirring under nitrogen for 16 h. The reaction mixture was diluted with 5% citric acid and was extracted with EtOAc. The organics were washed with water, saturated sodium bicarbonate and brine, dried over sodium sulfate and filtered. The product, VI, was purified by elution through silica in ethyl acetate and hexane.

Hydrolysis to final product: The methyl ester VI was dissolved in 1:1 tetrahydrofuran and methanol. With stirring was added aqueous lithium hydroxide (2 N). After stirring for one hour, the reaction mixture was concentrated to dryness. The residue was partitioned between 1 N aqueous hydrogen chloride and ethyl acetate, and the organic layer was washed with saturated sodium chloride. Drying over sodium sulfate, filtering and
25 evaporating gave crude acid. Purification by preparative reverse-phase high performance liquid chromatography gave pure acid VII.

The compounds of this invention may also be modified by appending appropriate functionalities to enhance selective biological properties. Such modifications are known in the art and include those which increase biological penetration into a given biological system
30 (e.g., blood, lymphatic system, central nervous system), increase oral availability, increase solubility to allow administration by injection, alter metabolism and alter rate of excretion.

Examples of these modifications include, but are not limited to, esterification with polyethylene glycols, derivatization with pivulates or fatty acid substituents, conversion to carbamates, hydroxylation of aromatic rings, and heteroatom-substitution in aromatic rings.

Also included are non-classical isoteres such as CO_2H , SO_2NHR , SO_3H ,



Once synthesized, the activities and VLA-4 specificities of the compounds according to this invention may be determined using *in vitro* and *in vivo* assays.

For example, the cell adhesion inhibitory activity of these compounds may be measured by determining the concentration of inhibitor required to block the binding of
 10 VLA-4-expressing cells to fibronectin- or CS1-coated plates. In this assay microtiter wells are coated with either fibronectin (containing the CS-1 sequence) or CS-1. If CS-1 is used, it must be conjugated to a carrier protein, such as bovine serum albumin, in order to bind to the wells. Once the wells are coated, varying concentrations of the test compound are then added together with appropriately labelled, VLA-4-expressing cells. Alternatively, the test
 15 compound may be added first and allowed to incubate with the coated wells prior to the addition of the cells. The cells are allowed to incubate in the wells for at least 30 minutes. Following incubation, the wells are emptied and washed. Inhibition of binding is measured by quantitating the fluorescence or radioactivity bound to the plate for each of the various concentrations of test compound, as well as for controls containing no test compound.

20 VLA-4-expressing cells that may be utilized in this assay include Ramos cells, Jurkat cells, A375 melanoma cells, as well as human peripheral blood lymphocytes (PBLs). The cells used in this assay may be fluorescently or radioactively labelled.

A direct binding assay may also be employed to quantitate the inhibitory activity of the compounds of this invention. In this assay, a VCAM-IgG fusion protein containing the
 25 first two immunoglobulin domains of VCAM (D1D2) attached above the hinge region of an IgG1 molecule ("VCAM 2D-IgG"), is conjugated to a marker enzyme, such as alkaline phosphatase ("AP"). The synthesis of this VCAM-IgG fusion is described in PCT publication WO 90/13300, the disclosure of which is herein incorporated by reference. The

conjugation of that fusion to a marker enzyme is achieved by cross-linking methods well-known in the art.

The VCAM-IgG enzyme conjugate is then placed in the wells of a multi-well filtration plate, such as that contained in the Millipore Multiscreen Assay System (Millipore Corp., Bedford, MA). Varying concentrations of the test inhibitory compound are then added to the wells followed by addition of VLA-4-expressing cells. The cells, compound and VCAM-IgG enzyme conjugate are mixed together and allowed to incubate at room temperature.

Following incubation, the wells are vacuum drained, leaving behind the cells and any bound VCAM. Quantitation of bound VCAM is determined by adding an appropriate colorimetric substrate for the enzyme conjugated to VCAM-IgG and determining the amount of reaction product. Decreased reaction product indicates increased binding inhibitory activity.

In order to assess the VLA-4 inhibitory specificity of the compounds of this invention, assays for other major groups of integrins, i.e., $\beta 2$ and $\beta 3$, as well as other $\beta 1$ integrins, such as VLA-5, VLA-6 and $\alpha 4\beta 7$ are performed. These assays may be similar to the adhesion inhibition and direct binding assays described above, substituting the appropriate integrin-expressing cell and corresponding ligand. For example, polymorphonuclear cells (PMNs) express $\beta 2$ integrins on their surface and bind to ICAM. $\beta 3$ integrins are involved in platelet aggregation and inhibition may be measured in a standard platelet aggregation assay. VLA-5 binds specifically to Arg-Gly-Asp sequences, while VLA-6 binds to laminin. $\alpha 4\beta 7$ is a recently discovered homologue of VLA-4, which also binds fibronectin and VCAM. Specificity with respect to $\alpha 4\beta 7$ is determined in a binding assay that utilizes the above-described VCAM-IgG-enzyme marker conjugate and a cell line that expresses $\alpha 4\beta 7$, but not VLA-4, such as RPMI-8866 cells.

Once VLA-4-specific inhibitors are identified, they may be further characterized in in vivo assays. One such assay tests the inhibition of contact hypersensitivity in an animal, such as described by P.L. Chisholm et al., "Monoclonal Antibodies to the Integrin $\alpha 4$ Subunit Inhibit the Murine Contact Hypersensitivity Response", Eur. J. Immunol., 23, pp. 682-688 (1993) and in "Current Protocols in Immunology", J. E. Coligan, et al., Eds., John Wiley & Sons, New York, 1, pp. 4.2.1-4.2.5 (1991), the disclosures of which is herein

incorporated by reference. In this assay, the skin of the animal is sensitized by exposure to an irritant, such as dinitrofluorobenzene, followed by light physical irritation, such as scratching the skin lightly with a sharp edge. Following a recovery period, the animals are re-sensitized following the same procedure. Several days after sensitization, one ear of the
5 animal is exposed to the chemical irritant, while the other ear is treated with a non-irritant control solution. Shortly after treating the ears, the animals are given various doses of the VLA-4 inhibitor by subcutaneous injection. In vivo inhibition of cell adhesion-associated inflammation is assessed by measuring the ear swelling response of the animal in the treated versus untreated ear. Swelling is measured using calipers or other suitable instrument to
10 measure ear thickness. In this manner, one may identify those inhibitors of this invention which are best suited for inhibiting inflammation.

Another in vivo assay that may be employed to test the inhibitors of this invention is the sheep asthma assay. This assay is performed essentially as described in W. M. Abraham et al., "α-Integrins Mediate Antigen-induced Late Bronchial Responses and Prolonged
15 Airway Hyperresponsiveness in Sheep", J. Clin. Invest., 93, pp. 776-87 (1994), the disclosure of which is herein incorporated by reference. This assay measures inhibition of *Ascaris* antigen-induced late phase airway responses and airway hyperresponsiveness in asthmatic sheep.

The compounds of the present invention may be used in the form of pharmaceutically
20 acceptable salts derived from inorganic or organic acids and bases. Included among such acid salts are the following: acetate, adipate, alginate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, citrate, camphorate, camphorsulfonate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, fumarate, glucoheptanoate, glycerophosphate, hemisulfate, heptanoate, hexanoate, hydrochloride, hydrobromide, hydroiodide, 2-
25 hydroxyethanesulfonate, lactate, maleate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, oxalate, pamoate, pectinate, persulfate, 3-phenyl-propionate, picrate, pivalate, propionate, succinate, tartrate, thiocyanate, tosylate and undecanoate. Base salts include ammonium salts, alkali metal salts, such as sodium and potassium salts, alkaline earth metal salts, such as calcium and magnesium salts, salts with organic bases, such as
30 dicyclohexylamine salts, N-methyl-D-glucamine, and salts with amino acids such as arginine, lysine, and so forth. Also, the basic nitrogen-containing groups can be quaternized

with such agents as lower alkyl halides, such as methyl, ethyl, propyl, and butyl chloride, bromides and iodides; dialkyl sulfates, such as dimethyl, diethyl, dibutyl and diamyl sulfates, long chain halides such as decyl, lauryl, myristyl and stearyl chlorides, bromides and iodides, aralkyl halides, such as benzyl and phenethyl bromides and others. Water or oil-soluble or dispersible products are thereby obtained.

The compounds of the present invention may be formulated into pharmaceutical compositions that may be administered orally, parenterally, by inhalation spray, topically, rectally, nasally, buccally, vaginally or via an implanted reservoir. The term "parenteral" as used herein includes subcutaneous, intravenous, intramuscular, intra-articular, intra-synovial, intrasternal, intrathecal, intrahepatic, intralesional and intracranial injection or infusion techniques.

The pharmaceutical compositions of this invention comprise any of the compounds of the present invention, or pharmaceutically acceptable derivatives thereof, together with any pharmaceutically acceptable carrier. The term "carrier" as used herein includes acceptable adjuvants and vehicles. Pharmaceutically acceptable carriers that may be used in the pharmaceutical compositions of this invention include, but are not limited to, ion exchangers, alumina, aluminum stearate, lecithin, serum proteins, such as human serum albumin, buffer substances such as phosphates, glycine, sorbic acid, potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts or electrolytes, such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, cellulose-based substances, polyethylene glycol, sodium carboxymethylcellulose, polyacrylates, waxes, polyethylene-polyoxypropylene-block polymers, polyethylene glycol and wool fat.

According to this invention, the pharmaceutical compositions may be in the form of a sterile injectable preparation, for example a sterile injectable aqueous or oleaginous suspension. This suspension may be formulated according to techniques known in the art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally

employed as a solvent or suspending medium. For this purpose, any bland fixed oil may be employed including synthetic mono- or di-glycerides. Fatty acids, such as oleic acid and its glyceride derivatives are useful in the preparation of injectables, as do natural pharmaceutically-acceptable oils, such as olive oil or castor oil, especially in their polyoxyethylated versions. These oil solutions or suspensions may also contain a long-chain alcohol diluent or dispersant, such as Ph. Helv or similar alcohol.

The pharmaceutical compositions of this invention may be orally administered in any orally acceptable dosage form including, but not limited to, capsules, tablets, aqueous suspensions or solutions.

In the case of tablets for oral use, carriers which are commonly used include lactose and corn starch. Lubricating agents, such as magnesium stearate, are also typically added. For oral administration in a capsule form, useful diluents include lactose and dried corn starch. When aqueous suspensions are required for oral use, the active ingredient is combined with emulsifying and suspending agents. If desired, certain sweetening, flavoring or coloring agents may also be added.

Alternatively, the pharmaceutical compositions of this invention may be administered in the form of suppositories for rectal administration. These can be prepared by mixing the agent with a suitable non-irritating excipient which is solid at room temperature but liquid at the rectal temperature and therefore will melt in the rectum to release the drug. Such materials include cocoa butter, beeswax and polyethylene glycols.

The pharmaceutical compositions of this invention may also be administered topically, especially when the target of treatment includes areas or organs readily accessible by topical application, including diseases of the eye, the skin, or the lower intestinal tract. Suitable topical formulations are readily prepared for each of these areas or organs.

Topical application for the lower intestinal tract can be effected in a rectal suppository formulation (see above) or in a suitable enema formulation. Topically-transdermal patches may also be used.

For topical applications, the pharmaceutical compositions may be formulated in a suitable ointment containing the active component suspended or dissolved in one or more carriers. Carriers for topical administration of the compounds of this invention include, but are not limited to, mineral oil, liquid petrolatum, white petrolatum, propylene glycol,

polyoxyethylene, polyoxypropylene compound, emulsifying wax and water. Alternatively, the pharmaceutical compositions can be formulated in a suitable lotion or cream containing the active components suspended or dissolved in one or more pharmaceutically acceptable carriers. Suitable carriers include, but are not limited to, mineral oil, sorbitan monostearate, polysorbate 60, cetyl esters wax, cetearyl alcohol, 2-octyldodecanol, benzyl alcohol and water.

For ophthalmic use, the pharmaceutical compositions may be formulated as micronized suspensions in isotonic, pH adjusted sterile saline, or, preferably, as solutions in isotonic, pH adjusted sterile saline, either with or without a preservative such as benzylalkonium chloride. Alternatively, for ophthalmic uses, the pharmaceutical compositions may be formulated in an ointment such as petrolatum.

The pharmaceutical compositions of this invention may also be administered by nasal aerosol or inhalation through the use of a nebulizer, a dry powder inhaler or a metered dose inhaler. Such compositions are prepared according to techniques well-known in the art of pharmaceutical formulation and may be prepared as solutions in saline, employing benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability, fluorocarbons, and/or other conventional solubilizing or dispersing agents.

The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated, and the particular mode of administration. It should be understood, however, that a specific dosage and treatment regimen for any particular patient will depend upon a variety of factors, including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, rate of excretion, drug combination, and the judgment of the treating physician and the severity of the particular disease being treated. The amount of active ingredient may also depend upon the therapeutic or prophylactic agent, if any, with which the ingredient is co-administered.

As stated above, an effective amount of a pharmaceutical composition containing an effective amount of a compound of this invention is also within the scope of this invention. An effective amount is defined as the amount which is required to confer a therapeutic effect on the treated patient, and will depend on a variety of factors, such as the nature of the inhibitor, the size of the patient, the goal of the treatment, the nature of the pathology to be

treated, the specific pharmaceutical composition used, and the judgment of the treating physician. For reference, see Freireich et al., Cancer Chemother. Rep. 1966, 50, 219 and Scientific Tables, Geigy Pharmaceuticals, Ardley, New York, 1970, 537. Dosage levels of between about 0.001 and about 100 mg/kg body weight per day, preferably between about 0.1 and about 10 mg/kg body weight per day of the active ingredient compound are useful.

According to another embodiment compositions containing a compound of this invention may also comprise an additional agent selected from the group consisting of corticosteroids, bronchodilators, antiasthmatics (mast cell stabilizers), antiinflammatories, antirheumatics, immunosuppressants, antimetabolites, immunomodulators, antipsoriatics and antidiabetics. Specific compounds within each of these classes may be selected from any of those listed under the appropriate group headings in "Comprehensive Medicinal Chemistry", Pergamon Press, Oxford, England, pp. 970-986 (1990), the disclosure of which is herein incorporated by reference. Also included within this group are compounds such as theophylline, sulfasalazine and aminosalicylates (antiinflammatories); cyclosporin, FK-506, and rapamycin (immunosuppressants); cyclophosphamide and methotrexate (antimetabolites); and interferons (immunomodulators).

According to other embodiments, the invention provides methods for preventing, inhibiting or suppressing cell adhesion-associated inflammation and cell adhesion-associated immune or autoimmune responses. VLA4-associated cell adhesion plays a central role in a variety of inflammation, immune and autoimmune diseases. Thus, inhibition of cell adhesion by the compounds of this invention may be utilized in methods of treating or preventing inflammatory, immune and autoimmune diseases. Preferably the diseases to be treated with the methods of this invention are selected from asthma, arthritis, psoriasis, transplantation rejection, multiple sclerosis, diabetes and inflammatory bowel disease.

These methods may employ the compounds of this invention in a monotherapy or in combination with an anti-inflammatory or immunosuppressive agent. Such combination therapies include administration of the agents in a single dosage form or in multiple dosage forms administered at the same time or at different times.

In order that this invention may be more fully understood, the following examples are set forth. These examples are for the purpose of illustration only and are not to be construed as limiting the scope of the invention in any way.

Intermediate 1:

4-(2-methylphenylaminocarbonylamino)phenylacetic Acid (oMePUPA-OH): To a suspension of *p*-aminophenylacetic acid (56.8 g, 376 mmol) in DMS (150 mL) was added *o*-tolyl isocyanate (50 g, 376 mmol) dropwise. The reaction mixture was allowed to stir 1 h, and was poured into EtOAc (1.75 L) with stirring. The precipitate was collected and washed with EtOAc (400 mL) and MeCN (400 mL) to provide oMePUPA (80 g, 75%). ESMS m/z ($M+H^+$) 285.1.

10 Intermediate 2:

OMePUPA-Leu-OH: oMePUPA-OH (0.78 g) was combined with Leucine methyl ester hydrochloride (0.50 g, 1.0 eq), HATU (1.10 g, 1.05 eq), and diisopropylethylamine (1.9 mL, 4 eq) in 10 mL dry DMF. The reaction was stirred for 16 h at room temperature after which it was diluted with 50 mL EtOAc, which was washed with 5% citric acid, water, saturated sodium bicarbonate and brine. The resulting organic solution was dried over sodium sulfate filtered and concentrated to yield 1.13 g of white solid. This product was dissolved in 10 mL THF. 5 mL 2N LiOH was added and the reaction was stirred for 16 h. THF was removed under reduced pressure and the solution was diluted with 40 mL water and washed with EtOAc. The aqueous layer was acidified with 1N HCl and was extracted with EtOAc. The organic extracts were washed with dilute HCl and brine, were dried over sodium sulfate, filtered and concentrated under reduced pressure yielding 0.77 g of white solid. ESMS m/z ($M+H^+$) 398.5.

Intermediate 3:

25 *N*-(3,5-dichlorobenzenesulfonyl)-Proline Methyl Ester: To a solution of 24.8 g (0.15 mol) of L-Proline methyl ester hydrochloride in 500 mL of CH_2Cl_2 was added 70 mL (0.5 mol) of triethylamine with stirring to give copious white precipitate. The mixture was filtered, and the filtrate cooled to 0° C (ice bath) with stirring. To the cooled solution was added a solution of 36.8 g (0.15 mol) of 3,5-dichlorobenzenesulfonyl chloride in 70 mL of CH_2Cl_2 dropwise quickly over five minutes. The addition funnel was rinsed with an additional 30 mL of CH_2Cl_2 , and the cloudy yellow mixture was allowed to warm to room temperature

with stirring overnight. The mixture was washed 2x with 400mL of 1N HCl, 2x with 400mL of 1N NaOH, then brine, then dried (MgSO₄), filtered, and concentrated to a yellow oil which crystallized on standing. The material was recrystallized three times from ethyl acetate/hexanes to give 39.3 g (0.116 mol, 77%) of *N*-(3,5-dichlorobenzenesulfonyl)-Proline methyl ester (MW = 338) as white needles (TLC on silica vs. 2:1 hexanes/ethyl acetate, R_f = 0.51). M/z = 339.3 (M+H⁺).

N-(3,5-diChlorobenzenesulfonyl)-Proline: To a solution of 39.3 g (0.116 mol) of the above methyl ester in 250 mL methanol was added 115 mL (0.23 mol) of freshly-prepared 2M aqueous LiOH with stirring to give a colorless solution. This was stirred for three hours, after which HPLC showed no starting material. The solution was reduced by 50% in vacuo and partitioned between 1N HCl and CH₂Cl₂ (~200 mL each). The phases were separated and the aqueous layer was washed again with CH₂Cl₂. The organic phases were combined, dried (MgSO₄), and concentrated to a white, foamy solid. This was recrystallized twice from ethyl acetate/hexanes to give 33.8 g (0.104 mol, 90%) of the title compound as colorless, broad, flat needles. M/z = 325.2 (M+H⁺).

Intermediate 4:




N-(benzenesulfonyl)-Proline Methyl Ester: To a solution of 25 g (0.15 mol) of L-Proline methyl ester hydrochloride in 500mL of CH₂Cl₂ was added 70 mL (0.5 mol) of triethylamine with stirring to give copious white precipitate. The mixture was filtered and the filtrate cooled to 0° C (ice bath) with stirring. To the cooled solution was added a solution of 20 mL (0.15 mol) of benzenesulfonyl chloride in 50 mL of CH₂Cl₂ dropwise over fifteen minutes. The addition funnel was rinsed with an additional 25 mL of CH₂Cl₂, and the cloudy, colorless mixture was allowed to warm to room temperature with stirring overnight. The solution was washed 2x with 400mL of 1N HCl, 2x with 400mL of 1N NaOH, 1x with brine, then dried (MgSO₄), filtered, and concentrated to a pale yellow solid. This material was recrystallized three times from ethyl acetate/hexanes to give 38.2 g (0.142 mol, 95%) of *N*-(benzenesulfonyl)-Proline methyl ester (MW = 269) as broad white needles (TLC vs. 2:1 hexanes/ethyl acetate, R_f = 0.35). M/z = 270.2 (M+H⁺).




N-(benzenesulfonyl)-Proline: To a solution of 38.2 g (0.142 mol) of the above methyl ester in 500 mL methanol was added 140 mL (0.28 mol) of freshly-prepared 2M aqueous LiOH


with stirring to give a colorless solution. This was stirred overnight, after which HPLC showed no starting material. The solution was reduced by 50% in vacuo and partitioned between 1N HCl and CH₂Cl₂ (~200 mL each). The phases were separated and the aqueous layer was washed again with CH₂Cl₂. The organic phases were combined, dried (MgSO₄),
5 and concentrated to a white solid. This was recrystallized twice from ethyl acetate/hexanes to give 34.7 g (0.136 mol, 96%) of the title compound as fine white needles. M/z = 256.2 (M+H⁺).

Example 1

10 Synthesis of Compound IX







Methyl ester Hydrochloride I: In a 500 mL RB flask was suspended 8.4 g (33.3 mmol) 2-N-CBZ-L-2,4-diaminobutyric acid in 200 mL methanol (MeOH) with stirring. This was cooled to 0 degrees C (ice bath), and then 14.6 mL (200 mmol) SOCl_2 was added dropwise over 15 minutes to give a colorless solution. The solution was allowed to warm to RT and stirred overnight, after which a proton NMR spectrum of an aliquot indicated the

reaction was complete. The solution was concentrated, redissolved in MeOH and concentrated 2x, then dissolved in CH₂Cl₂, conc., and placed under high vacuum for 16 hours to give compound I as a slightly yellow foam, massing to 10.33g (34.2 mmol, 103%). MS: m/z 267 (M+H)⁺.

- 5 tert-Butoxycarbonyl methyl ester II: In a 500mL RB flask was dissolved 10.33 g (33.3 mmol) of I in dry dimethylformamide (DMF) with stirring to give a colorless solution. To this was added 17.4 mL (100 mmol) of diisopropylethylamine (DIEA), then 7.96 g (32.5 mmol) of Boc-N-Methyl-Leucine, and finally 14.83 g (39.0 mmol) of O-(7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (HATU) to give a yellow solution.
- 10 This was stirred overnight, after which HPLC showed no starting material. The solution was diluted with ethyl acetate (EtOAc, 500mL) and washed with 1N HCl (2x), 1N NaOH (2x), and brine (1x). The organic phase was dried over anhydrous MgSO₄, filtered, and concentrated to a red oil. Chromatography with 2:1 hexanes/EtOAc vs. silica gave 12.56 g (25.5 mmol, 78%) of II as a yellow syrup (HPLC, >99%). MS: m/z 393 (M-BOC)⁺, 494
- 15 (M+H)⁺.

- Amino ester III: In a 280 mL high-pressure vessel was dissolved 11.38 g (23.08 mmol) of II in 75 mL MeOH with stirring to give an orange solution. The vessel was flushed with nitrogen, and ~200mg (catalytic) of 10% palladium on carbon (Pd/C) was added. The sides of the vessel were washed with additional MeOH, and the vessel capped with a
- 20 hydrogenation head. The mixture was placed under 60 psi H₂ with stirring overnight, after which HPLC showed no starting material remained. The mixture was filtered through Celite 545, the filter pad rinsed with additional MeOH, and the filtrate concentrated to a colorless oil, III, massing to 8.29 g (~quantitative). Material carried through. MS: m/z 360 (M+H)⁺.

- Benzyl carbamate methyl ester IV: In a 500 mL RB flask was dissolved 8.29 g
- 25 (23.08 mmol) of III in 100mL CH₂Cl₂ with stirring to give a colorless solution. To this was added 7.0 mL (50 mmol) of triethylamine (Et₃N), then 7.96 g (23.0 mmol) of CBZ-proline hydroxysuccinimide ester (CBZ-Pro-Osu) to give a colorless solution. This was stirred overnight, after which HPLC showed no starting material remaining. The solution was diluted with additional CH₂Cl₂, washed with 1N HCl (2x), 1N NaOH (2x), and the organic
- 30 phase dried over MgSO₄, filtered, and the filtrate concentrated to a colorless oil.

Chromatography with 3:1 EtOAc/hexanes vs. silica gave 12.22 g (20.7 mmol, 90%) of IV as a foamy, colorless glass (HPLC, >99%). MS: m/z 490 (M-BOC)⁺, 591 (M+H)⁺.

Amine trifluoroacetate salt V: In a 500 mL RB flask was dissolved 11.80 g (20.0 mmol) of IV in 120 mL CH₂Cl₂ with stirring to give a colorless solution. To this was added
5 20 mL (260 mmol, large excess) of trifluoroacetic acid (TFA), and the resulting solution was stirred for four hours, after which HPLC showed no starting material. The solution was concentrated, redissolved in CH₂Cl₂ and concentrated (2x), then placed under high vacuum to give 12.1 g (~quantitative) of V as a pale yellow oil. Material carried through. MS: m/z 491 (M+H)⁺.


10 Diaryl urea methyl ester VI: In a 500 mL RB flask was dissolved 12.1 g (20 mmol) of V in 100 mL DMF with stirring to give a pale yellow solution. To this was added 17.4 mL (100 mmol) of DIEA, then 5.68 g (20.0 mmol) Intermediate 1 (oMePUPA-OH), and finally 9.12 g (24 mmol) of HATU to give a yellow solution. This was stirred overnight, after which HPLC showed no starting material remaining. The solution was diluted with EtOAc (500
15 mL) and washed with 1N HCl (2x), 1N NaOH (2x), and brine (1x). The organic phase was dried with MgSO₄, filtered, and the filtrate concentrated to a yellow oil/solid mixture. Chromatography with 2:1 acetonitrile/CH₂Cl₂ vs. silica gave 11.35 g (15.0 mmol, 75%) of VI as a slightly yellow, foamy solid (HPLC, >99%). MS: m/z 757 (M+H)⁺, 779 (M+Na⁺).

Amino methyl ester VII: In a 280 mL high-pressure vessel was dissolved 8.0 g (10.6
20 mmol) of VI in 50 mL MeOH with stirring to give a slightly yellow solution. The vessel was flushed with nitrogen, and ~250 mg (catalytic) of 10% Pd/C added. The sides of the vessel were washed with additional MeOH and the vessel capped with the hydrogenation head. The mixture was placed under 60 psi H₂ with stirring overnight, after which HPLC showed no starting material. The mixture was filtered through Celite 545, the filter pad rinsed with
25 additional MeOH, and the filtrate concentrated to give 6.6 g (~quantitative) of VII as a white solid. Material carried through. MS: m/z 623 (M+H)⁺.

Sulfonamide methyl ester VIII: In a 500 mL RB flask was dissolved 6.6 g (10.6 mmol) of VII in 100 mL dry CH₂Cl₂ with stirring to give a colorless solution. This was cooled to 0 degrees C (ice bath), and 4.2 mL (30 mmol) of Et₃N was added, followed by a solution of 3.68
30 g (15 mmol) of 3,5-dichlorobenzenesulfonyl chloride in 25 mL dry CH₂Cl₂ added dropwise over 10 minutes. The resulting solution was allowed to warm to RT and stirred for 2 hours,

after which HPLC showed no starting material. The solution was diluted with additional CH_2Cl_2 and washed with 1N HCl (2x) and 1N NaOH (2x), then dried over MgSO_4 , filtered, and the filtrate concentrated to a yellow solid. Chromatography with 2:1 CH_2Cl_2 /acetonitrile vs. silica gave 6.68 g (8.0 mmol, 75%) of VIII as a white solid (HPLC, >99%). MS: m/z 832/833 ($\text{M}+\text{H}^+$).

Carboxylic acid IX: In a 500 mL RB flask was dissolved 6.26 g (7.53 mmol) of VIII in 150 mL MeOH with stirring to give a colorless solution. This was cooled to 0 degrees C (ice bath), and nitrogen was bubbled through the stirring solution for 30 minutes. To this was added 19 mL (38 mmol) of freshly-made 2M LiOH solution dropwise over 10 minutes, after which the solution was stirred at 0 degrees C under nitrogen while the reaction progress was closely monitored by HPLC. After three hours, HPLC showed no starting material remaining. The solution was concentrated with minimal heating (volume reduced ~ 50%), and slowly poured, in portions, into ice-cold 1N HCl to give a copious, brilliant-white precipitate. The solid was isolated via filtration, washed with cold distilled water, and air-dried overnight. The resulting fine, white solid was transferred to a glass jar and placed under high vacuum for 72 hours. The final mass was 6.02 g (7.36 mmol, 98%) of IX as a white powder (HPLC, >98%). MS: m/z 818/819 ($\text{M}+\text{H}^+$), 841 ($\text{M}+\text{Na}^+$).

Example 2:**Synthesis of Compound XVI**

5

Homoserine 4-nitrophenyl Ether Benzyl Ester: To a solution of N-Boc homoserine benzyl ester **I** (1.2 g, 3.89 mmol), 4-nitrophenol (485 mg, 4.08 mmol) and triphenylphosphine (1.2 g, 4.66 mmol) in THF (10 mL) diethylazodicarboxylate (DEAD) (0.74 mL, 4.66 mmol) was added dropwise and the reaction was stirred at room temperature 12-24h. Upon completion as judged by LC the solvents were removed to afford a viscous syrup. 4N HCl in dioxane (10 mL) was added rapidly and the solution was stirred at room temperature 3-6 h or until judged complete by LC. The reaction was concentrated to ¼

10

volume and the product was precipitated out of ethyl acetate to afford the hydrochloride salt II (96% pure, LC) as a white solid (867 mg, 2.36 mmol, 61%). ESMS: (M-Cl) = 331.

To a solution of Intermediate 4 (117 mg, 0.46 mmol) in DMF (3 mL) was added DIPEA (0.27 mL, 1.84 mmol) followed sequentially by the hydrochloride salt II (160 mg, 0.48 mmol) and HATU (239 mg, 0.63 mmol). The solution was stirred at room temperature for 2-4 h until judged complete by LC. The reaction was diluted with ethyl acetate (30 mL) and washed with 5% bicarbonate (10 mL), water (10 mL), citric acid (10 mL), brine (2 x 10 mL) and dried over sodium sulfate to afford the crude product III as a tan foam (213 mg, 0.37 mmol, 82%) which was used directly.

ESMS: (M+H) = 568.

The above material was dissolved in ethyl acetate (15 mL), 10% Pd/C (200 mg) was added and the reaction was subjected to hydrogenolysis at 50 psi for 4-6 h or until judged complete by LC. Filtration through celite and concentration afforded the crude aniline IV (144 mg, 0.32 mmol, 87%) as a tan foam which was used immediately.













ESMS: (M+H) = 448.











The aniline (74 mg, 0.17 mmol) obtained above was dissolved in DMF (3 mL) and oMePUPA (52 mg, 0.18 mmol) was added followed by DIPEA (0.08 mL, 0.43 mmol) and HATU (69 mg, 0.18 mmol) and the reaction was stirred at room temperature 3-4 h until complete by LC. Purification by HPLC afforded Bio-8355 (39 mg, 0.054 mmol, 30%) as a white solid.













ESMS: (M+H) = 714, (M-H) = 712.

Compounds of this invention as shown in the following tables were prepared according to the method described above.









Compounds prepared according to General Method A include:







Compound #	R3	R1	ESMS m/z
5450			610.7 (M+H ⁺)
5451			589.3 (M+H ⁺)
6668			498.2 (M+H ⁺)
6669			468.1 (M+H ⁺)
6670			534.5 (M+H ⁺)
6671			484.4 (M+H ⁺)










6697	oMePUPA-Pro		774.3 (M+H ⁺)
6714	oMePUPA-N-MeLeu		804.4 (M+H ⁺)
6715			670 (M+H ⁺)
6716			686.4 (M+H ⁺)
7171			505.2 (M+H ⁺)
7172			475.2 (M+H ⁺)











7175			541.3 (M+H ⁺)
7177			491.6 (M+H ⁺)
7514			678.3 (M+H ⁺)
7515			662.4 (M+H ⁺)
7516			692.3 (M+H ⁺)
7517			676.6 (M+H ⁺)






Compounds prepared according to General Method B include:



BIO#	R3	R1	ESMS m/z
7855	oMePUPCH2		664.3 (M+H ⁺)
7856			560.2 (M+H ⁺)
7857			532.1 (M+H ⁺)
8066	CH3		440.0 (M+H ⁺)
8067	Bn		516.0 (M+H ⁺)
8122	oMePUPCH2		539.5 (M+H ⁺)

8123			435.4 (M+H ⁺)
8147			419.0 (M+H ⁺)
8208	oMePUPCH2	CH3	469.0 (M+H ⁺)
8209	oMePUPCH2	oMePUPCH2	693.1 (M+H ⁺)
8210		CH3	507.9 (M+H ⁺)
8211		oMePUPCH2	732.3 (M+H ⁺)










8212			771.1 (M+H ⁺)
8449	oMePUPCH2		573.0 (M+H ⁺)
8450	Bn		425.0 (M+H ⁺)
8451			557.9 (M+H ⁺)
8452			469.0 (M+H ⁺)
8453	oMePUPCH2		600.0 (M+H ⁺)









8455			585.0 (M+H ⁺)
8456			495.9 (M+H ⁺)
8457			546.0 (M+Na ⁺)
8458	oMePUPCH2		745.9 (M+H ⁺)
8459	Bn		597.9 (M+H ⁺)
8460			730.9 (M+H ⁺)










8461			641.8 (M+H ⁺)
8462	oMePUPCH2	oMePUPA-Leu	806.1 (M+H ⁺)
8463	Bn	oMePUPA-Leu	658.1 (M+H ⁺)
8464		oMePUPA-Leu	791.0 (M+H ⁺)
8465		CH3	454.0 (M+H ⁺)
8466		CH3	365.0 (M+H ⁺)





8519	 <chem>CC(F)(F)c1ccccc1</chem>	 <chem>CC1CCCN1S(=O)(=O)c2cc(Cl)cc(Cl)c2</chem>	633.8 (M+H ⁺)
------	--------------------------------------------------------------------------------------------------------------------	-------------------------------------------------------------------------------------------------------------------------------------	---------------------------

Compounds prepared according to General Method C include:







Compound #	R3	R1	ESMS m/z
5801			518.0(M+H ⁺)
5803	oMePUPCH2		650.0 (M+H ⁺)
6655		CH3	344.2 (M+H ⁺)
7081			546.0 (M+H ⁺)
7111			659.7 (M+H ⁺)
7117		CH3	351.2 (M+H ⁺)







7119	oMePUPCH2	CH3	452.8 (M+H ⁺)
7147			602.2 (M+H ⁺)
7148			539.1 (M+H ⁺)
7150	2-Cl-Bn		642.1 (M+H ⁺)
7156	oMePUPCH2		740.2 (M+H ⁺)
7157			636.1 (M+H ⁺)







7158	CH ₃		516.2 (M+H ⁺)
7231	H		452.1 (M+H ⁺)
7233			616.1 (M+H ⁺)
7234	oMePUPA-Leu		831.1 (M+H ⁺)
7235			642.0 (M+H ⁺)
7236			639.0 (M+H ⁺)







7241	oMePUPCH2		664.3 (M+H ⁺)
7255	PhCH2CO-Pro		667.1 (M+H ⁺)
7256	oMePUPA-Pro		815.1 (M+H ⁺)
7257	PhCH2CO-Leu		683.1 (M+H ⁺)







Compounds prepared according to General Method D include:






Compound #	R1	ESMS m/z
5292		620.8 (M-H ⁺)
7080		743.9 (M+H ⁺)
7092		875.8 (M+H ⁺)
7093		843.8 (M+H ⁺)
7109		843.8 (M+H ⁺)
7116		905.7 (M+H ⁺)



7181		833.1 (M+H ⁺)
7200		713.4 (M+H ⁺)
7328		685.0 (M-H ⁺)
7398		832.1 (M+H ⁺)
7662		750.1 (M+H ⁺)
8221		832.9 (M+H ⁺)

8290		703.1 (M+H ⁺)
8291		703.1 (M+H ⁺)
8294		720.1 (M+H ⁺)
8295		720.1 (M+H ⁺)
8308		741.1 (M+H ⁺)
8309		803.1 (M+H ⁺)











8341		750.0 (M+H ⁺)
8493		765.9 (M+H ⁺)
8528		966.1 (M+H ⁺)
8555		764.0 (M+H ⁺)
8571		735.2 (M+H ⁺)
8582		826.0 (M+H ⁺)













8583		764.1 (M+H ⁺)
8586		791.1 (M+H ⁺)
8628		763.2 (M+H ⁺)
8642		754.0 (M+H ⁺)
8674		764.1 (M+H ⁺)
8929		686.2 (M+H ⁺)















9120		852.2 (M+H ⁺)
9140	_CH3	554.2 (M+H ⁺)
9169		881.4 (M+H ⁺)
9170		783.3 (M+H ⁺)
9171		791.3 (M+H ⁺)
9182		775.5 (M+H ⁺)







9264		764.2 (M+H ⁺)
9437		903.3 (M+H ⁺)

Compounds prepared according to General Method E include:

Compound #	R3	L	R1	ESMS m/z
5800	Ac-Leu-			824.7 (M+H ⁺)
7083	oMePUPCH2			850.5 (M+H ⁺)
7155	oMePUPCH2	-(CH2)3-		705.9 (M+H ⁺)
7168	PhCH2CO-N-Me-Leu	-(CH2)2-		565.2(M+H ⁺)
7528		-(CH2)2-		691.0 (M+H ⁺)
7530		-(CH2)2-		675.0 (M+H ⁺)

7552	oMePUPA- α -N-Me- ϵ -CBz-Lys-	-(CH ₂) ₂ -		968.1 (M+H ⁺)
7578	oMePUPA-N-Me-Gly	-(CH ₂) ₂ -		785.0 (M+Na ⁺)
9232	oMePUPCH ₂			770.2 (M-H ⁺)
9233	oMePUPA-Leu			883.6 (M-H ⁺)
9234	oMePUPCH ₂			625.1 (M+H ⁺)
9235	oMePUPA-Leu			738.2 (M+H ⁺)
9236	oMePUPCH ₂			786.2 (M+H ⁺)

9237	oMePUPA-Leu			897.4 (M-H ⁺)
9238	oMePUPCH2			639.1 (M+H ⁺)
9239	oMePUPA-Leu			750.1 (M-H ⁺)
9270	oMePUPCH2			742.1 (M-H ⁺)
9271	oMePUPA-Leu			855.4 (M-H ⁺)
9273	oMePUPA-Leu			710.1 (M+H ⁺)
9274	oMePUPCH2			758.1 (M+H ⁺)

9275	oMePUPA-Leu			869.2 (M+H ⁺)
9276	oMePUPCH2			611.0 (M+H ⁺)
9277	oMePUPA-Leu			724.1 (M+H ⁺)

Other Embodiments

From the above description, one skilled in the art can easily ascertain the essential characteristics of the present invention, and without departing from the spirit and scope thereof, can make various changes and modifications of the invention to adapt it to various usages and conditions. Thus, other embodiments are also within the claims.

WHAT IS CLAIMED IS:

1 1. A compound of the formula:



3 wherein

4 R^1 is

- 5 1) H,
- 6 2) C_{1-10} alkyl,
- 7 3) C_{2-10} alkenyl,
- 8 4) C_{2-10} alkynyl,
- 9 5) Cy,
- 10 6) Cy- C_{1-10} alkyl,
- 11 7) Cy- C_{1-10} alkenyl, or
- 12 8) Cy- C_{1-10} alkynyl;

13 L' is a hydrocarbon linker moiety having 1-5 carbon chain atoms and is

14 (i) optionally interrupted by, or terminally attached to, one or more of the following
15 groups:

- 16 1) -C(O)-,
- 17 2) -O-C(O)-,
- 18 3) -C(O)-O-,
- 19 4) -C(O)-NR^c-,
- 20 5) -NR^c-C(O)-,
- 21 6) -NR^c-C(O)-NR^d-,
- 22 7) -NR^c-C(O)-O-,
- 23 8) -O-C(O)-NR^c-,
- 24 9) -S(O)_m-,
- 25 10) -SO₂-NR^c-,
- 26 11) -NR^c-SO₂-,
- 27 12) -NR^c-C(NR^m)-,
- 28 13) -O-,
- 29 14) -NR^c-, or
- 30 15) -Cy; or

31 (ii) optionally substituted with one or more substituents independently selected from R^b;

32 L is a hydrocarbon linker moiety having 1-14 carbon chain atoms and is


33 (i) optionally interrupted by, or terminally attached to, one or more of the following
34 groups:

- 35 1) -C(O)-,
- 36 2) -O-C(O)-,
- 37 3) -C(O)-O-,
- 38 4) -C(O)-NR^c-,
- 39 5) -NR^c-C(O)-,
- 40 6) -NR^c-C(O)-NR^d-,
- 41 7) -NR^c-C(O)-O-,
- 42 8) -O-C(O)-NR^c-,
- 43 9) -S(O)_m-,
- 44 10) -SO₂-NR^c-,
- 45 11) -NR^c-SO₂-,
- 46 12) -O-,
- 47 13) -NR^c-, or
- 48 14) Cy; or

49 (ii) optionally substituted with one or more substituents independently selected from R^b;

50 and

51 R³ is alkyl, alkenyl, alkynyl, cycloalkyl, aryl-fused cycloalkyl, cycloalkenyl, aryl, aralkyl,
52 aryl-substituted alkenyl or alkynyl, cycloalkyl-substituted alkyl, cycloalkenyl-substituted
53 cycloalkyl, biaryl, alkenoxy, alkynoxy, aralkoxy, aryl-substituted alkenoxy or alkynoxy,
54 alkylamino, alkenylamino or alkynylamino, aryl-substituted alkylamino, aryl-substituted
55 alkenylamino or alkynylamino, aryloxy, arylamino, heterocyclyl, heterocyclyl-substituted
56 alkyl, heterocyclyl-substituted amino, carboxyalkyl substituted aralkyl, or oxocarbocyclyl-
57 fused aryl; or a moiety of the following formula:



58

59 wherein:

60 Y^5 is selected from the group consisting of -CO-, -O-CO-, -SO₂- and -PO₂-;
61 each of R^4 and R^6 , independently, is alkyl, alkenyl, alkynyl, cycloalkyl, aryl-fused
62 cycloalkyl, cycloalkenyl, aryl, aralkyl, aryl-substituted alkenyl or alkynyl, cycloalkyl-
63 substituted alkyl, cycloalkenyl-substituted cycloalkyl, biaryl, alkenoxy, alkynoxy, aralkoxy,
64 aryl-substituted alkenoxy or alkynoxy, alkylamino, alkenylamino or alkynylamino, aryl-
65 substituted alkylamino, aryl-substituted alkenylamino or alkynylamino, aryloxy, arylamino,
66 heterocyclyl, heterocyclyl-substituted alkyl, heterocyclyl-substituted amino, carboxyalkyl
67 substituted aralkyl, oxocarbocyclyl-fused aryl, or an amino acid side chain selected from the
68 group consisting of arginine, asparagine, glutamine, S-methyl cysteine, methionine and
69 corresponding sulfoxide and sulfone derivatives thereof, cyclohexylalanine, leucine,
70 isoleucine, allo-isoleucine, tert-leucine, norleucine, phenylalanine, phenylglycine, tyrosine,
71 tryptophan, proline, alanine, ornithine, histidine, glutamine, norvaline, valine, threonine,
72 serine, beta-cyanoalanine, 2-aminobutyric acid and allothreonine; and
73 R^5 is hydrogen, aryl, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, or aryl-substituted
74 alkyl, R^5 and R^6 may be taken together with the atoms to which they are attached to form a
75 heterocycle of 5 to 7 members;

76 each of said Cy is cycloalkyl, cycloalkenyl, heterocyclyl, aryl, or heteroaryl;

77 each of said alkyl, alkenyl and alkynyl is optionally substituted with one to four
78 substituents independently selected from R^a ; and

79 each of said cycloalkyl, cycloalkenyl, heterocyclyl, aryl, or heteroaryl is optionally
80 substituted with one to four substituents independently selected from R^b ;

81 R^a is

- 82 1) Cy,
- 83 2) -OR^c,
- 84 3) -NO₂,
- 85 4) -halogen,
- 86 5) -S(O)_mR^c,
- 87 6) -SR^c,
- 88 7) -S(O)₂OR^c,
- 89 8) -S(O)₂NR^cR^d,

- 90 9) $-\text{NR}^{\text{c}}\text{R}^{\text{d}}$,
- 91 10) $-\text{O}(\text{CR}^{\text{c}}\text{R}^{\text{f}})_n\text{NR}^{\text{c}}\text{R}^{\text{d}}$,
- 92 11) $-\text{C}(\text{O})\text{R}^{\text{d}}$,
- 93 12) $-\text{CO}_2\text{R}^{\text{c}}$,
- 94 13) $-\text{P}(\text{O})(\text{OR}^{\text{c}})(\text{OR}^{\text{d}})$,
- 95 14) $-\text{P}(\text{O})(\text{R}^{\text{c}})(\text{OR}^{\text{d}})$,
- 96 15) $-\text{S}(\text{O})_m\text{OR}^{\text{c}}$,
- 97 16) $-\text{C}(\text{O})\text{NR}^{\text{c}}\text{R}^{\text{j}}$,
- 98 17) $-\text{CO}_2(\text{CR}^{\text{c}}\text{R}^{\text{f}})_n\text{CONR}^{\text{c}}\text{R}^{\text{d}}$,
- 99 18) $-\text{OC}(\text{O})\text{R}^{\text{c}}$,
- 100 19) $-\text{CN}$,
- 101 20) $-\text{NR}^{\text{c}}\text{C}(\text{O})\text{R}^{\text{d}}$,
- 102 21) $-\text{OC}(\text{O})\text{NR}^{\text{c}}\text{R}^{\text{d}}$,
- 103 22) $-\text{NR}^{\text{c}}\text{C}(\text{O})\text{OR}^{\text{d}}$,
- 104 23) $-\text{NR}^{\text{c}}\text{C}(\text{O})\text{NR}^{\text{d}}\text{R}^{\text{e}}$,
- 105 24) $-\text{CR}^{\text{c}}(\text{NOR}^{\text{d}})$,
- 106 25) $-\text{CF}_3$,
- 107 26) $-\text{OCF}_3$, or
- 108 27) oxo

109 wherein Cy is optionally substituted with one to four substituents independently selected
 110 from R^{b} ,

111 R^{b} is

- 112 1) a group selected from R^{a} ,
- 113 2) C_{1-10} alkyl,
- 114 3) C_{2-10} alkenyl,
- 115 4) C_{2-10} alkynyl,
- 116 5) aryl- C_{1-10} alkyl, or
- 117 6) heteroaryl- C_{1-10} alkyl,

118 wherein each of alkyl, alkenyl, alkynyl, aryl, and heteroaryl is optionally substituted with a
 119 group independently selected from R^{g}

120 each of R^{c} , R^{d} , R^{e} , and R^{f} , independently, is

- 121 1) H,
122 2) C₁₋₁₀ alkyl,
123 3) C₂₋₁₀ alkenyl,
124 4) C₂₋₁₀ alkynyl,
125 5) Cy, or
126 6) Cy-C₁₋₁₀ alkyl;

127 wherein each of alkyl, alkenyl, alkynyl and Cy is optionally substituted with one to four
128 substituents independently selected from R^g;

129 R^g is

- 130 1) halogen,
131 2) amino,
132 3) carboxy,
133 4) -COO-C₁₋₄ alkyl,
134 5) -P(O)(OH)₂,
135 6) -P(O)(OH)(O-C₁₋₄ alkyl),
136 7) -P(O)(C₁₋₄ alkyl)₂,
137 8) -P(O)(OH)(C₁₋₄ alkyl),
138 9) -P(O)(O-C₁₋₄ alkyl)(C₁₋₄ alkyl),
139 10) -SO₂-C₁₋₄ alkyl,
140 11) -CO-NH₂,
141 12) -CO-NH(C₁₋₄ alkyl),
142 13) -CO-N(C₁₋₄ alkyl)₂,
143 14) C₁₋₄ alkyl,
144 15) C₁₋₄ alkoxy,
145 16) aryl,
146 17) aryl-C₁₋₄ alkoxy,
147 18) hydroxy,
148 19) CF₃, or
149 20) aryloxy;

150 R^m is

- 151 1) H,

- 152 2) C₁₋₁₀ alkyl,
153 3) C₂₋₁₀ alkenyl,
154 4) C₂₋₁₀ alkynyl,
155 5) Cy,
156 6) Cy-C₁₋₁₀ alkyl;
157 7) C₁₋₁₀ acyl,
158 8) cyano,
159 9) C₁₋₁₀ alkyl-sulfonyl, or
160 10) C₁₋₁₀ alkoxy; and

161 R^j is

- 162 1) H,
163 2) C₁₋₁₀ alkyl,
164 3) C₂₋₁₀ alkenyl,
165 4) C₂₋₁₀ alkynyl,
166 5) cyano,
167 6) aryl,
168 7) aryl-C₁₋₁₀ alkyl,
169 8) heteroaryl,
170 9) heteroaryl-C₁₋₁₀ alkyl, or
171 10) -SO₂R^k,

172 where R^k is C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl, and aryl;

173 R^c and R^d taken together with the atoms to which they are attached optionally form a
174 heterocyclic ring of 5 to 7 members, said ring containing 0-2 additional heteroatoms
175 independently selected from O, N and S;

176 R^e and R^f taken together with the atoms to which they are attached optionally form a
177 ring of 5 to 7 members, said ring containing 0-2 additional heteroatoms independently
178 selected from O, S and N;

179 m is 0, 1, or 2;

180 n is an integer from 1 to 10;

181 provided that when L is saturated and has 1-4 carbon chain atoms,

182 (i) L must contain a heteroatom selected from O, S, and N; or

- 183 (ii) R^3 must contain the moiety o-methylphenyl-ureido-phenyl- CH_2 -; or
184 (iii) R^1 must contain only one Cy group;
185 or a pharmaceutically acceptable salt thereof.

186

1 2. The compound of claim 1, wherein R^1 is $Z^1-L^a-Z^2$ -,

2 in which

3 Z^1 is cycloalkyl, cycloalkyl- C_{1-10} alkyl, cycloalkenyl, cycloalkenyl- C_{1-10} alkyl, aryl, aryl-
4 C_{1-10} alkyl, heterocyclyl, heterocyclyl- C_{1-10} alkyl, heteroaryl, or heteroaryl- C_{1-10} alkyl;

5 L^a is -C(O)-, -O-C(O)-, -C(O)-O-, -C(O)- NR^c -, - NR^c -C(O)-, - NR^c -C(O)- NR^d -, - NR^c -
6 C(O)-O-, -O-C(O)- NR^c -, -S(O) $_m$ -, -SO $_2$ - NR^c -, - NR^c -SO $_2$ -, -O-, - NR^c -, or a bond; m, R^c and
7 R^d having been defined in claim 1; and

8 Z^2 is cycloalkyl, cycloalkyl- C_{1-10} alkyl, cycloalkenyl, cycloalkenyl- C_{1-10} alkyl, aryl, aryl-
9 C_{1-10} alkyl, heterocyclyl, heterocyclyl- C_{1-10} alkyl, heteroaryl, heteroaryl- C_{1-10} alkyl or a bond.

1

1 3. The compound of claim 2, wherein

2 Z^1 is cycloalkyl, cycloalkyl- C_{1-10} alkyl, aryl, aryl- C_{1-10} alkyl, heterocyclyl, heterocyclyl-
3 C_{1-10} alkyl, heteroaryl, or heteroaryl- C_{1-10} alkyl;

4 L^a is -O-C(O)-, -C(O)-O-, -C(O)- NR^c -, - NR^c -C(O)-, -SO $_2$ -, -SO $_2$ - NR^c -, - NR^c -SO $_2$ -, -O-, -
5 NR^c -, or a bond; and

6 Z^2 is aryl, aryl- C_{1-10} alkyl, heterocyclyl, heterocyclyl- C_{1-10} alkyl, or a bond.

1 4. The compound of claim 3, wherein

2 Z^1 is aryl, aryl- C_{1-5} alkyl, heterocyclyl, heterocyclyl- C_{1-5} alkyl, heteroaryl, or heteroaryl-
3 C_{1-5} alkyl;

4 L^a is -O-C(O)-, -C(O)-O-, -C(O)- NR^c -, - NR^c -C(O)-, -SO $_2$ -, or a bond; and

5 Z^2 is heterocyclyl, heterocyclyl- C_{1-5} alkyl, or a bond.


1 5. The compound of claim 4, wherein

2 Z^1 is phenyl optionally substituted with Cy, -CO- R^d , halogen, oxo, aryl-substituted
3 alkenyl;

4 L^a is -O-C(O)-, -C(O)-O-, -C(O)- NR^c -, - NR^c -C(O)-, or -SO $_2$ -; and

5 Z^2 is heterocyclyl or a bond.

1 6. The compound of claim 1, wherein R^1 is



2 R^9 is

- 3
- 4 1) C_{1-10} alkyl,
 - 5 2) C_{2-10} alkenyl,
 - 6 3) C_{2-10} alkynyl,
 - 7 4) Cy,
 - 8 5) Cy- C_{1-10} alkyl,
 - 9 6) Cy- C_{2-10} alkenyl, or
 - 10 7) Cy- C_{2-10} alkynyl;

11 each of R^{10} and R^{11} , independently, is selected from the group consisting of hydrogen, aryl,
12 alkyl, alkenyl or alkynyl, cycloalkyl, cycloalkenyl, and aryl-substituted alkyl;


13 R^{12} is

- 14
- 15 1) H,
 - 16 2) C_{1-10} alkyl,
 - 17 3) C_{2-10} alkenyl,
 - 18 4) C_{2-10} alkynyl,
 - 19 5) aryl,
 - 20 6) aryl- C_{1-10} alkyl,
 - 21 7) heteroaryl, or
 - 22 8) heteroaryl- C_{1-10} alkyl;

23 wherein each of alkyl, alkenyl and alkynyl is optionally substituted with one to four
24 substituents independently selected from R^a , and aryl and heteroaryl are optionally
25 substituted with one to four substituents independently selected from R^b ;

26 R^{11} , R^{12} and the carbon to which they are attached form a 3-7 membered mono- or bicyclic
ring containing 0-2 heteroatoms selected from N, O, and S.

1 7. The compound of claim 1, wherein R^1 is



2

3

R^{14} is

4

8) C_{1-10} alkyl,

5

9) C_{2-10} alkenyl,

6

10) C_{2-10} alkynyl,

7

11) Cy,

8

12) Cy- C_{1-10} alkyl,

9

13) Cy- C_{2-10} alkenyl, or

10

14) Cy- C_{2-10} alkynyl,

11

R^{15} is

12

1) H,

13

2) C_{1-10} alkyl,

14

3) C_{2-10} alkenyl,

15

4) C_{2-10} alkynyl,

16

5) aryl,

17

6) aryl- C_{1-10} alkyl,

18

7) heteroaryl, or

19

8) heteroaryl- C_{1-10} alkyl,

20

each of R^{16} , R^{17} , and R^{18} , independently, is

21

1) H,

22

2) C_{1-10} alkyl,

23

3) C_{2-10} alkenyl,

24

4) C_{2-10} alkynyl,

25

5) Cy,

26


6) Cy- C_{1-10} alkyl,

- 27 7) Cy-C₂₋₁₀ alkenyl,
 28 8) Cy-C₂₋₁₀ alkynyl, or
 29 9) a group selected from R^a

30 wherein Cy is optionally substituted with one to four substituents independently selected
 31 from R^b or one of the following groups:

- 32 1) -NR^cC(O)NR^cSO₂R^d,
 33 2) -NR^cS(O)_mR^d,
 34 3) -OS(O)₂OR^c, or
 35 4) -OP(O)(OR^c)₂;

36 two of R¹⁶, R¹⁷, and R¹⁸, when attached to a common ring atom, together with the common
 37 ring atom form a 5-7 membered saturated or unsaturated monocyclic ring containing zero to
 38 three heteroatoms selected from N, O, or S; or two of R¹⁶, R¹⁷, and R¹⁸, when attached to two
 39 adjacent ring atoms, together with these two ring atoms form a 5-7 membered saturated or
 40 unsaturated monocyclic ring containing zero to three heteroatoms selected from N, O, or S;



41 the ring represents a 3-7 membered saturated or unsaturated heterocycl
 42 or heteroaryl wherein

43 each of Z, A, B₁ and B₂, independently, is

- 44 1) a bond,
 45 2) -C-,
 46 3) -C-C-,
 47 4) -C=C-,
 48 5) a heteroatom selected from the group consisting of N, O, and S, or
 49 6) -S(O)_m-;

50 m having been defined in claim 1;


51 Y⁷ is


- 52 1) -C(O)-,
 53 2) -C(O)O-,
 54 3) -C(O)NR^c-,
 55 4) -S(O)₂-,


56 5) $-P(O)(OR^c)$, or

57 6) $-C(O)-C(O)-$;

58 wherein each of said alkyl, alkenyl and alkynyl is optionally substituted with one to four
 59 substituents independently selected from R^a , and each said Cy is optionally substituted with
 60 one to four substituents independently selected from R^b .

1 8. The compound of claim 7, wherein the ring  represents azetidine,
 2 pyrrole, pyrrolidine, imidazole, pyrazole, triazole, pyridine, piperidine, pyrazine, piperazine,
 3 pyrimidine, oxazole, thiazole, or morpholine.

1 9. The compound of claim 8, wherein the ring  represents azetidine,
 2 pyrrole, pyrrolidine, imidazole, piperidine, or morpholine.

1 10. The compound of claim 9, wherein the ring  represents pyrrolidine.


1 11. The compound of claim 7, wherein R^{15} is H or C_{1-5} alkyl.

1 12. The compound of claim 7, wherein each of R^{16} , R^{17} , and R^{18} , independently, is
 2 selected from the group consisting of H, C_{1-10} alkyl, Cy, $-OR^c$, -halogen, $-S(O)_mR^c$, $-NR^cR^d$, -
 3 $NR^cC(O)R^d$, $-NR^cC(O)OR^d$, $-NR^cC(O)NR^dR^e$, and oxo; each of R^c , R^d , R^e , and m having
 4 been defined in claim 1.

1 13. The compound of claim 7, wherein Y^7 is $-O-C(O)-$, $-C(O)-O-$, or $-SO_2-$.

1 14. The compound of claim 13, wherein Y^7 is $-SO_2-$.

- 1 15. The compound of claim 7, wherein R^{14} is Cy or Cy- C_{1-5} alkyl.
- 1 16. The compound of claim 15, wherein Cy is phenyl.
- 1 17. The compound of claim 1, wherein L' contains 2-4 carbon chain atoms.
- 1 18. The compound of claim 1, wherein L' is



1 in which

2 Y^1 is

- 3 15) $-C(O)-$,
- 4 16) $-O-C(O)-$,
- 5 17) $-C(O)-O-$,
- 6 18) $-C(O)-NR^c-$,
- 7 19) $-NR^c-C(O)-$,
- 8 20) $-NR^c-C(O)-NR^d-$,
- 9 21) $-NR^c-C(O)-O-$,
- 10 22) $-O-C(O)-NR^c-$,
- 11 23) $-S(O)_m-$,
- 12 24) $-S(O)_2-NR^c-$,
- 13 25) $-NR^c-S(O)_2-$,
- 14 26) $-NR^c-C(NR^m)-$,
- 15 27) $-O-$, or
- 16 28) $-NR^c-$;

17 R^2 is

- 18 1) H,
- 19 2) C_{1-10} alkyl,

- 21 3) C₂₋₁₀ alkenyl,
- 22 4) C₂₋₁₀ alkynyl,
- 23 5) Cy,
- 24 6) Cy-C₁₋₁₀ alkyl,
- 25 7) Cy-C₁₋₁₀ alkenyl, or
- 26 8) Cy-C₁₋₁₀ alkynyl;

27 Y² is a bond or -C(R^h)(Rⁱ)-; wherein

28 each of R^h and Rⁱ is independently selected from the group consisting of:

- 29 1) H,
- 30 2) C₁₋₁₀ alkyl,
- 31 3) C₂₋₁₀ alkenyl,
- 32 4) C₂₋₁₀ alkynyl,
- 33 5) aryl,
- 34 6) aryl-C₁₋₁₀ alkyl,
- 35 7) heteroaryl, and
- 36 8) heteroaryl-C₁₋₁₀ alkyl,

37 R^h and Rⁱ taken together with the carbon to which they are attached may optionally form a 3-
38 7 membered ring containing 0-2 heteroatoms selected from N, O and S;

39 X is

- 40 1) -C(O)OR^c,
- 41 2) -P(O)(OR^c)(OR^d),
- 42 3) -P(O)(R^c)(OR^d),
- 43 4) -S(O)_mOR^c,
- 44 5) -C(O)NR^cR^j, or
- 45 6) -5-tetrazolyl;

46 m having been defined in claim 1;

47 wherein each of said alkyl, alkenyl and alkynyl is optionally substituted with one to four
48 substituents independently selected from R^a, each of said aryl and heteroaryl is optionally
49 substituted with one to four substituents independently selected from R^b; and Cy is a
50 cycloalkyl, heterocyclyl, aryl, or heteroaryl; and
51 provided that when Y² is not a bond, X is -COOH, -COO-C₁₋₄ alkyl, -P(O)(OH)₂,

52 -P(O)(OH)(O-C₁₋₄ alkyl), -P(O)(C₁₋₄ alkyl)₂, -P(O)(OH)(C₁₋₄ alkyl), -P(O)(O-C₁₋₄ alkyl)(C₁₋₄
 53 alkyl), -SO₂-C₁₋₄ alkyl, -CO-NH₂, -CO-NH(C₁₋₄ alkyl), -CO-N(C₁₋₄ alkyl)₂, or -5-tetrazolyl.

1 19. The compound of claim 18, wherein Y¹ is -NR^c-C(O)-, -NR^c-, -NR^c-S(O)₂-, or -NR^c-
 2 C(NR^m)₂-.

1 20. The compound of claim 19, wherein Y¹ is -NR^c-C(O)-.

1 21. The compound of claim 18, wherein R² is H or C₁₋₅ alkyl.

1 22. The compound of claim 21, wherein R² is H.

1 23. The compound of claim 18, wherein Y² is a bond or -C(R^h)(Rⁱ)-, wherein each of R^h
 2 and Rⁱ, independently, is H or C₁₋₅ alkyl.

1 24. The compound of claim 23, wherein each of R^h and Rⁱ, independently, is H.

1 25. The compound of claim 23, wherein Y² is a bond.


1 26. The compound of claim 18, wherein X is -C(O)OR^c or -C(O)NR^cR^j.

1 27. The compound of claim 26, wherein X is -C(O)OR^c where R^c is H or C₁₋₅ alkyl.

1 28. The compound of claim 18, wherein Y¹ is -NR^c-C(O)-; R² is H or C₁₋₅ alkyl; Y² is a
 2 bond or -CH₂-; and X is -C(O)OR^c where each R^c is independently H or C₁₋₅ alkyl.

1 29. The compound of claim 1, wherein L contains 4-10 carbon chain atoms.

1 30. The compound of claim 1, wherein L is



2 in which

3 Y³ is

- 5 9) a bond,
- 6 10) C₁₋₁₀ alkyl,
- 7 11) C₂₋₁₀ alkenyl,
- 8 12) C₂₋₁₀ alkynyl,
- 9 13) aryl,
- 10 14) aryl-C₁₋₁₀ alkyl,
- 11 15) heteroaryl, or
- 12 16) heteroaryl-C₁₋₁₀ alkyl; and

13 Y⁴ is

- 14 1) a bond,
- 15 2) -C(O)-,
- 16 3) -O-C(O)-,
- 17 4) -C(O)-O-,
- 18 5) -C(O)-NR^c-,
- 19 6) -NR^c-C(O)-,
- 20 7) -NR^c-C(O)-NR^d-,
- 21 8) -NR^c-C(O)-O-,
- 22 9) -O-C(O)-NR^c-,
- 23 17) -S(O)_m-,
- 24 18) -S(O)₂-NR^c-,
- 25 19) -NR^c-S(O)₂-,
- 26 20) -NR^c-C(NR^d)-,
- 27 21) -O-, or
- 28 22) -NR^c;

29 wherein each of alkyl, alkenyl, and alkynyl is optionally containing one to four heteroatoms
 30 selected from N, O, S, and -S(O)_m-, and each of alkyl, alkenyl and alkynyl is optionally
 31 substituted with one to four substituents independently selected from R^a, and each of aryl and
 32 heteroaryl is optionally substituted with one to four substituents independently selected from
 33 R^b; each of R^a, R^b, R^c, R^d, and m having been defined in claim 1; and
 34 provided that each of Y³ and Y⁴ is not a bond simultaneously.

1 31. The compound of claim 30, wherein Y^3 is a bond, C_{1-5} alkyl, or C_{1-5} alkenyl; and Y^4
 2 is a bond, $-C(O)-NR^c$ -, $-C(O)-$ -, $-NR^c$ -, or $-O-$, where R^c is H or C_{1-5} alkyl.

1 32. The compound of claim 1, wherein

2 R^3 is $Z^3-L^b-Z^4$ -, in which

3 Z^3 is Cy, Cy- C_{1-10} alkyl, Cy- C_{1-10} alkenyl, or Cy- C_{1-10} alkynyl;

4 L^b is $-C(O)-$ -, $-O-C(O)-$ -, $-C(O)-O-$ -, $-C(O)-NR^c$ -, $-NR^c-C(O)-$ -, $-NR^c-C(O)-NR^d$ -, $-NR^c$ -


5 $C(O)-O-$ -, $-O-C(O)-NR^c$ -, $-S(O)_m$ -, $-SO_2-NR^c$ -, $-NR^c-SO_2$ -, $-O-$ -, $-NR^c$ -, or a bond; and

6 Z^4 is cycloalkyl, cycloalkyl- C_{1-10} alkyl, cycloalkenyl, cycloalkenyl- C_{1-10} alkyl, aryl, aryl-

7 C_{1-10} alkyl, heterocyclyl, heterocyclyl- C_{1-10} alkyl, heteroaryl, heteroaryl- C_{1-10} alkyl or a

8 bond; or

9 R^3 is a moiety of the formula:



10

11 each of m , R^c , R^d , R^4 , R^5 , R^6 , and Y^5 having been defined in claim 1.

1 33. The compound of claim 32, wherein R^4 is $Z^5-L^c-Z^6$ -,

2 in which

3 Z^5 is Cy, Cy- C_{1-10} alkyl, Cy- C_{1-10} alkenyl, or Cy- C_{1-10} alkynyl;

4 L^c is $-C(O)-$ -, $-O-C(O)-$ -, $-C(O)-O-$ -, $-C(O)-NR^c$ -, $-NR^c-C(O)-$ -, $-NR^c-C(O)-NR^d$ -, $-NR^c$ -

5 $C(O)-O-$ -, $-O-C(O)-NR^c$ -, $-S(O)_m$ -, $-SO_2-NR^c$ -, $-NR^c-SO_2$ -, $-O-$ -, $-NR^c$ -, or a bond; and

6 Z^6 is cycloalkyl, cycloalkyl- C_{1-10} alkyl, cycloalkenyl, cycloalkenyl- C_{1-10} alkyl, aryl, aryl-

7 C_{1-10} alkyl, heterocyclyl, heterocyclyl- C_{1-10} alkyl, heteroaryl, heteroaryl- C_{1-10} alkyl or a

8 bond;

9 each of m , R^c , and R^d having been defined in claim 1.

1 34. The compound of claim 33, wherein

2 each of Z^3 and Z^5 , independently, is aryl, aryl- C_{1-10} alkyl, aryl- C_{1-10} alkenyl, aryl- C_{1-10}
3 alkynyl, heteroaryl, heteroaryl- C_{1-10} alkyl, heteroaryl- C_{1-10} alkenyl, or heteroaryl- C_{1-10}
4 alkynyl;
5 each of L^b and L^c , independently, is $-C(O)-$, $-S(O)_m-$, $-O-C(O)-$, $-C(O)-O-$, $-C(O)-NR^c-$, $-$
6 $NR^c-C(O)-$, $-NR^c-C(O)-NR^d-$, $-SO_2-NR^c-$, $-NR^c-SO_2-$, $-O-$, $-NR^c-$, or a bond; and
7 each of Z^4 and Z^6 , independently, is aryl, aryl- C_{1-10} alkyl, heterocyclyl, heterocyclyl- C_{1-10}
8 alkyl, heteroaryl, heteroaryl- C_{1-10} alkyl, or a bond.

1 35. The compound of claim 34, wherein
2 each of Z^3 and Z^5 , independently, is aryl, aryl- C_{1-10} alkyl, heteroaryl, or heteroaryl- C_{1-10}
3 alkyl;
4 each of L^b and L^c , independently, is $-C(O)-$, $-SO_2-$, $-C(O)-NR^c-$, $-NR^c-C(O)-$, or $-NR^c-$
5 $C(O)-NR^d-$; where each of R^c and R^d , independently, is H or C_{1-5} alkyl; and
6 each of Z^4 and Z^6 , independently, is aryl, aryl- C_{1-10} alkyl, heterocyclyl, heterocyclyl- C_{1-10}
7 alkyl, heteroaryl, heteroaryl- C_{1-10} alkyl, or a bond.

1 36. The compound of claim 35, wherein
2 each of Z^3 and Z^5 , independently, is aryl;
3 each of L^b and L^c , independently, is $-NR^c-C(O)-NR^d-$; and
4 each of Z^4 and Z^6 , independently, is aryl.

1 37. The compound of claim 32, wherein Y^5 is $-CO-$ or $-O-CO-$.

1 38. The compound of claim 37, wherein Y^5 is $-CO-$.

1 39. The compound of claim 32, wherein R^5 is H or C_{1-5} alkyl.

1 40. The compound of claim 39, wherein R^5 is H or C_{1-2} alkyl.

1 41. The compound of claim 32, wherein R^6 is an amino acid side chain selected from the
2 group consisting of cyclohexylalanine, leucine, isoleucine, allo-isoleucine, tert-leucine,

norleucine, phenylalanine, phenylglycine, alanine, norvaline, valine, and 2-aminobutyric acid.

42. The compound of claim 41, wherein R^6 is an amino acid side chain selected from the group consisting of leucine, isoleucine, allo-isoleucine, tert-leucine, norleucine, alanine, norvaline, valine, and 2-aminobutyric acid.

43. The compound of claim 42, wherein R^6 is the side chain of leucine or isoleucine.

44. The compound of claim 32, wherein R^1 is $Z^1-L^a-Z^2$,

in which


Z^1 is aryl optionally substituted with Cy, $-CO-R^d$, halogen, oxo, or aryl-substituted alkenyl;

L^a is $-O-C(O)-$, $-C(O)-O-$, $-C(O)-NR^c-$, $-NR^c-C(O)-$, or $-SO_2-$; and

Z^2 is heteroaryl, heterocyclyl, or a bond.

45. The compound of claim 44, wherein Z^1 is phenyl; L^a is $-SO_2-$; and Z^2 is azetidine, pyrrole, pyrrolidine, imidazole, piperidine, or morpholine.

46. The compound of claim 44, wherein L' is




in which

Y^1 is $-NR^c-C(O)-$, $-NR^c-$, $-NR^c-S(O)_2-$, or $-NR^c-C(NR^d)-$; R^2 is H or C_{1-5} alkyl; Y^2 is a bond or $-C(R^h)(R^i)-$; and X is $-C(O)OR^c$; where each of R^c , R^h , and R^i , independently, is H or C_{1-5} alkyl.

47. The compound of claim 46, wherein Y^1 is $-NH-C(O)-$; R^2 is H; Y^2 is a bond; and X is $-C(O)OH$.


- 1 48. The compound of claim 46, wherein L is



- 2
3 wherein Y³ is a bond, C₁₋₅ alkyl, or C₁₋₅ alkenyl; and Y⁴ is a bond, -C(O)-NR^c-, -C(O)-, -
4 NR^c-, or -O-, where R^c is H or C₁₋₅ alkyl.

- 1 49. The compound of claim 48, wherein Y³ is a bond or C₁₋₅ alkyl; and Y⁴ is -C(O)-NH-.

- 1 50. The compound of claim 48, wherein R³ is a moiety of the formula:



- 2
3 in which R⁴ is Z⁵-L^c-Z⁶-, where
4 Z⁵ is aryl, aryl-C₁₋₁₀ alkyl, aryl-C₁₋₁₀ alkenyl, aryl-C₁₋₁₀ alkynyl, heteroaryl, heteroaryl-C₁₋₁₀
5 alkyl, heteroaryl-C₁₋₁₀ alkenyl, or heteroaryl-C₁₋₁₀ alkynyl;
6 L^c is -C(O)-, -S(O)_m-, -O-C(O)-, -C(O)-O-, -C(O)-NR^c-, -NR^c-C(O)-, -NR^c-C(O)-NR^d-, -
7 SO₂-NR^c-, -NR^c-SO₂-, -O-, -NR^c-, or a bond, with R^c and R^d, independently, being H or
8 C₁₋₅ alkyl; and
9 Z⁶ is aryl, aryl-C₁₋₁₀ alkyl, heterocyclyl, heterocyclyl-C₁₋₁₀ alkyl, heteroaryl, heteroaryl-
10 C₁₋₁₀ alkyl, or a bond.

- 1 51. The compound of claim 50, wherein Z⁵ is aryl; L^c is -NR^c-C(O)-NR^d-; and Z⁶ is aryl.

- 1 52. The compound of claim 51, wherein R⁴ is o-methylphenyl-ureido-phenyl-CH₂-.


- 1 53. The compound of claim 51, wherein Y⁵ is -CO- or -O-CO-.

- 1 54. The compound of claim 53, wherein R⁵ is H or C₁₋₂ alkyl.


1 55. The compound of claim 54, wherein R^6 is an amino acid side chain selected from the
 2 group consisting of leucine, isoleucine, allo-isoleucine, tert-leucine, norleucine, alanine,
 3 norvaline, valine, and 2-aminobutyric acid.

1 56. The compound of claim 55, wherein R^6 is the side chain of leucine or isoleucine.

1 57. The compound of claim 1, wherein the chemical structure of said compound is



1 58. The compound of claim 1, wherein the chemical structure of said compound is



2
 3 wherein each of R^{21} and R^{22} , independently, is selected from a group consisting of

- 4 23) Cy,
- 5 24) $-OR^c$,
- 6 25) $-NO_2$,
- 7 26) -halogen,
- 8 27) $-S(O)_mR^c$,
- 9 28) $-SR^c$,
- 10 29) $-S(O)_2OR^c$,
- 11 30) $-S(O)_2NR^cR^d$,
- 12 31) $-NR^cR^d$,
- 13 32) $-O(CR^cR^f)_nNR^cR^d$,
- 14 33) $-C(O)R^c$,
- 15 34) $-CO_2R^c$,
- 16 35) $-CO_2(CR^cR^f)_nCONR^cR^d$,

- 17 36) -OC(O)R^c,
- 18 37) -CN,
- 19 38) -C(O)NR^cR^d,
- 20 39) -NR^cC(O)R^d,
- 21 40) -OC(O)NR^cR^d,
- 22 41) -NR^cC(O)OR^d,
- 23 42) -NR^cC(O)NR^dR^e,
- 24 43) -CR^c(NOR^d),
- 25 44) -CF₃,
- 26 45) -OCF₃,
- 27 46) oxo
- 28 47) C₁₋₁₀ alkyl,
- 29 48) C₂₋₁₀ alkenyl,
- 30 49) C₂₋₁₀ alkynyl,
- 31 50) aryl-C₁₋₁₀ alkyl, and
- 32 51) heteroaryl-C₁₋₁₀ alkyl

33 wherein each of alkyl, alkenyl, alkynyl, aryl, heteroaryl is optionally substituted with a
 34 group independently selected from R^e;

35 R²³ is selected from the group consisting of

- 36 1) H,
- 37 2) C₁₋₁₀ alkyl,
- 38 3) C₂₋₁₀ alkenyl,
- 39 4) C₂₋₁₀ alkynyl,
- 40 5) aryl,
- 41 6) aryl-C₁₋₁₀ alkyl,
- 42 7) heteroaryl, and
- 43 8) heteroaryl-C₁₋₁₀ alkyl,

44 wherein alkyl, alkenyl and alkynyl are optionally substituted with one to four substituents
 45 independently selected from R^a, and aryl and heteroaryl are optionally substituted with
 46 one to four substituents independently selected from R^b.

1 59. The compound of claim 1, where said compound is compound no. 5192, 5283, 6696,
 2 6697, 6714, 7234, 7256, 7578, 7662, 8221, 8308, 8309, 8341, 8342, 8343, 8367, 8368, 8469,
 3 8491, 8554, 8555, 8571, 8642, 8646, 8685, 8689, 8690, 8698, 8749, 8758, 8796, 8797, 8809,
 4 9120, 9169, 9171, 9182, 9227, 9264, 9271, 9315, 9418, 9621, 7083, 7200, 7328, 7399, 7788,
 5 7855, 8205, 8290, 8291, 8294, 8295, 8304, 8557, 8582, 8583, 8585, 8586, 8606, 8607, 8628,
 6 8674, 8684, 8723, 8746, 8929, 9273, or 9275.

1 60. The compound of claim 1, where said compound is compound nos. 7083, 7200, 7328,
 2 7399, 7788, 7855, 8205, 8290, 8291, 8294, 8295, 8304, 8557, 8582, 8583, 8585, 8586, 8606,
 3 8607, 8628, 8674, 8684, 8723, 8746, 8929, 9273, or 9275.

1 61. A composition comprising a pharmaceutical carrier and an effective amount of a
 2 compound of the following formula:



4 wherein

5 R^1 is

- 6 52) H,
- 7 53) C_{1-10} alkyl,
- 8 54) C_{2-10} alkenyl,
- 9 55) C_{2-10} alkynyl,
- 10 56) Cy,
- 11 57) Cy- C_{1-10} alkyl,
- 12 58) Cy- C_{1-10} alkenyl, or
- 13 59) Cy- C_{1-10} alkynyl;

14 L' is a hydrocarbon linker moiety having 1-5 carbon chain atoms and is

15 (i) optionally interrupted by, or terminally attached to, one or more of the following
 16 groups:

- 17 1) $-C(O)-$,
- 18 2) $-O-C(O)-$,
- 19 3) $-C(O)-O-$,
- 20 4) $-C(O)-NR^c-$,
- 21 5) $-NR^c-C(O)-$,

- 22 6) $-\text{NR}^c-\text{C}(\text{O})-\text{NR}^d-$,
23 7) $-\text{NR}^c-\text{C}(\text{O})-\text{O}-$,
24 8) $-\text{O}-\text{C}(\text{O})-\text{NR}^c-$,
25 9) $-\text{S}(\text{O})_m-$,
26 10) $-\text{SO}_2-\text{NR}^c-$,
27 11) $-\text{NR}^c-\text{SO}_2-$,
28 12) $-\text{NR}^c-\text{C}(\text{NR}^m)-$,
29 13) $-\text{O}-$,
30 14) $-\text{NR}^c-$, or
31 15) $-\text{Cy}$; or

32 (ii) optionally substituted with one or more substituents independently selected from R^b ;

33 L is a hydrocarbon linker moiety having 1-14 carbon chain atoms and is


34 (i) optionally interrupted by, or terminally attached to, one or more of the following
35 groups:

- 36 1) $-\text{C}(\text{O})-$,
37 2) $-\text{O}-\text{C}(\text{O})-$,
38 3) $-\text{C}(\text{O})-\text{O}-$,
39 4) $-\text{C}(\text{O})-\text{NR}^c-$,
40 5) $-\text{NR}^c-\text{C}(\text{O})-$,
41 6) $-\text{NR}^c-\text{C}(\text{O})-\text{NR}^d-$,
42 7) $-\text{NR}^c-\text{C}(\text{O})-\text{O}-$,
43 8) $-\text{O}-\text{C}(\text{O})-\text{NR}^c-$,
44 9) $-\text{S}(\text{O})_m-$,
45 10) $-\text{SO}_2-\text{NR}^c-$,
46 11) $-\text{NR}^c-\text{SO}_2-$,
47 12) $-\text{O}-$,
48 13) $-\text{NR}^c-$, or
49 14) Cy ; or

50 (ii) optionally substituted with one or more substituents independently selected from R^b ;

51 and

52 R^3 is alkyl, alkenyl, alkynyl, cycloalkyl, aryl-fused cycloalkyl, cycloalkenyl, aryl, aralkyl,
 53 aryl-substituted alkenyl or alkynyl, cycloalkyl-substituted alkyl, cycloalkenyl-substituted
 54 cycloalkyl, biaryl, alkenoxy, alkynoxy, aralkoxy, aryl-substituted alkenoxy or alkynoxy,
 55 alkylamino, alkenylamino or alkynylamino, aryl-substituted alkylamino, aryl-substituted
 56 alkenylamino or alkynylamino, aryloxy, arylamino, heterocyclyl, heterocyclyl-substituted
 57 alkyl, heterocyclyl-substituted amino, carboxyalkyl substituted aralkyl, or oxocarbocyclyl-
 58 fused aryl; or a moiety of the following formula:



59

60 wherein:

61 Y^5 is selected from the group consisting of $-\text{CO}-$, $-\text{O}-\text{CO}-$, $-\text{SO}_2-$ and $-\text{PO}_2-$;
 62 each of R^4 and R^6 , independently, is alkyl, alkenyl, alkynyl, cycloalkyl, aryl-fused
 63 cycloalkyl, cycloalkenyl, aryl, aralkyl, aryl-substituted alkenyl or alkynyl, cycloalkyl-
 64 substituted alkyl, cycloalkenyl-substituted cycloalkyl, biaryl, alkenoxy, alkynoxy, aralkoxy,
 65 aryl-substituted alkenoxy or alkynoxy, alkylamino, alkenylamino or alkynylamino, aryl-
 66 substituted alkylamino, aryl-substituted alkenylamino or alkynylamino, aryloxy, arylamino,
 67 heterocyclyl, heterocyclyl-substituted alkyl, heterocyclyl-substituted amino, carboxyalkyl
 68 substituted aralkyl, oxocarbocyclyl-fused aryl, or an amino acid side chain selected from the
 69 group consisting of arginine, asparagine, glutamine, S-methyl cysteine, methionine and
 70 corresponding sulfoxide and sulfone derivatives thereof, cyclohexylalanine, leucine,
 71 isoleucine, allo-isoleucine, tert-leucine, norleucine, phenylalanine, phenylglycine, tyrosine,
 72 tryptophan, proline, alanine, ornithine, histidine, glutamine, norvaline, valine, threonine,
 73 serine, beta-cyanoalanine, 2-aminobutyric acid and allothreonine; and
 74 R^5 is hydrogen, aryl, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, or aryl-substituted
 75 alkyl, R^5 and R^6 may be taken together with the atoms to which they are attached to form a
 76 heterocycle of 5 to 7 members;
 77 each of said Cy is cycloalkyl, heterocyclyl, aryl, or heteroaryl;
 78 each of said alkyl, alkenyl and alkynyl is optionally substituted with one to four
 79 substituents independently selected from R^a ; and

- 80 each of said cycloalkyl, cycloalkenyl, heterocyclyl, aryl, or heteroaryl is optionally
81 substituted with one to four substituents independently selected from R^b;
- 82 R^a is
- 83 1) Cy,
 - 84 2) -OR^c,
 - 85 3) -NO₂,
 - 86 4) -halogen,
 - 87 5) -S(O)_mR^c,
 - 88 6) -SR^c,
 - 89 7) -S(O)₂OR^c,
 - 90 8) -S(O)₂NR^cR^d,
 - 91 9) -NR^cR^d,
 - 92 10) -O(CR^cR^f)_nNR^cR^d,
 - 93 11) -C(O)R^d,
 - 94 12) -CO₂R^c,
 - 95 13) -P(O)(OR^c)(OR^d),
 - 96 14) -P(O)(R^c)(OR^d),
 - 97 15) -S(O)_mOR^c,
 - 98 16) -C(O)NR^cR^j,
 - 99 17) -CO₂(CR^cR^f)_nCONR^cR^d,
 - 100 18) -OC(O)R^c,
 - 101 19) -CN,
 - 102 20) -NR^cC(O)R^d,
 - 103 21) -OC(O)NR^cR^d,
 - 104 22) -NR^cC(O)OR^d,
 - 105 23) -NR^cC(O)NR^dR^e,
 - 106 24) -CR^c(NOR^d),
 - 107 25) -CF₃,
 - 108 26) -OCF₃, or
 - 109 27) oxo

110 wherein Cy is optionally substituted with one to four substituents independently selected
111 from R^b,

112 R^b is

- 113 1) a group selected from R^a,
- 114 2) C₁₋₁₀ alkyl,
- 115 3) C₂₋₁₀ alkenyl,
- 116 4) C₂₋₁₀ alkynyl,
- 117 5) aryl-C₁₋₁₀ alkyl, or
- 118 6) heteroaryl-C₁₋₁₀ alkyl,

119 wherein each of alkyl, alkenyl, alkynyl, aryl, and heteroaryl is optionally substituted with a
120 group independently selected from R^g

121 each of R^c, R^d, R^e, and R^f, independently, is

- 122 1) H,
- 123 2) C₁₋₁₀ alkyl,
- 124 3) C₂₋₁₀ alkenyl,
- 125 4) C₂₋₁₀ alkynyl,
- 126 5) Cy, or
- 127 6) Cy-C₁₋₁₀ alkyl;

128 wherein each of alkyl, alkenyl, alkynyl and Cy is optionally substituted with one to four
129 substituents independently selected from R^g,

130 R^g is

- 131 1) halogen,
- 132 2) amino,
- 133 3) carboxy,
- 134 4) -COO-C₁₋₄ alkyl,
- 135 5) -P(O)(OH)₂,
- 136 6) -P(O)(OH)(O-C₁₋₄ alkyl),
- 137 7) -P(O)(C₁₋₄ alkyl)₂,
- 138 8) -P(O)(OH)(C₁₋₄ alkyl),
- 139 9) -P(O)(O-C₁₋₄ alkyl)(C₁₋₄ alkyl),
- 140 10) -SO₂-C₁₋₄ alkyl,

- 141 11) -CO-NH₂,
142 12) -CO-NH(C₁₋₄ alkyl),
143 13) -CO-N(C₁₋₄ alkyl)₂,
144 14) C₁₋₄ alkyl,
145 15) C₁₋₄ alkoxy,
146 16) aryl,
147 17) aryl-C₁₋₄ alkoxy,
148 18) hydroxy,
149 19) CF₃, or
150 20) aryloxy;
151 R^m is
152 1) H,
153 2) C₁₋₁₀ alkyl,
154 3) C₂₋₁₀ alkenyl,
155 4) C₂₋₁₀ alkynyl,
156 5) Cy,
157 6) Cy-C₁₋₁₀ alkyl,
158 7) C₁₋₁₀ acyl,
159 8) C₁₋₁₀ alkyl-sulfonyl, or
160 9) C₁₋₁₀ alkoxy; and
161 R^j is
162 1) H,
163 2) C₁₋₁₀ alkyl,
164 3) C₂₋₁₀ alkenyl,
165 4) C₂₋₁₀ alkynyl,
166 5) cyano,
167 6) aryl,
168 7) aryl-C₁₋₁₀ alkyl,
169 8) heteroaryl,
170 9) heteroaryl-C₁₋₁₀ alkyl, or
171 10) -SO₂R^k,

172 where R^k is C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, or aryl;
 173 R^c and R^d taken together with the atoms to which they are attached optionally form a
 174 heterocyclic ring of 5 to 7 members, said ring containing 0-2 additional heteroatoms
 175 independently selected from O, N and S;
 176 R^e and R^f taken together with the atoms to which they are attached optionally form a
 177 ring of 5 to 7 members, said ring containing 0-2 additional heteroatoms independently
 178 selected from O, S and N;
 179 m is 0, 1, or 2;
 180 n is an integer from 1 to 10;
 181 provided that when L is saturated and has 1-4 carbon chain atoms,
 182 (i) L must contain a heteroatom selected from O, S, and N; or
 183 (ii) R^3 must contain the moiety o-methylphenyl-ureido-phenyl- CH_2 -; or
 184 (iii) R^1 must contain only one Cy group;
 185 or a pharmaceutically acceptable salt thereof.

1 62. The composition of claim 61, wherein said compound is compound nos. 5192, 5283,
 2 6696, 6697, 6714, 7234, 7256, 7578, 7662, 8221, 8308, 8309, 8341, 8342, 8343, 8367, 8368,
 3 8469, 8491, 8554, 8555, 8571, 8642, 8646, 8685, 8689, 8690, 8698, 8749, 8758, 8796, 8797,
 4 8809, 9120, 9169, 9171, 9182, 9227, 9264, 9271, 9315, 9418, 9621, 7083, 7200, 7328, 7399,
 5 7788, 7855, 8205, 8290, 8291, 8294, 8295, 8304, 8557, 8582, 8583, 8585, 8586, 8606, 8607,
 6 8628, 8674, 8684, 8723, 8746, 8929, 9273, or 9275

1 63. A method of inhibiting VLA-4-dependent cell adhesion, comprising administering to
 2 a patient in need thereof an effective amount of a compound of the following formula:



4 wherein

5 R^1 is

- 6 60) H,
- 7 61) C_{1-10} alkyl,
- 8 62) C_{2-10} alkenyl,
- 9 63) C_{2-10} alkynyl,
- 10 64) Cy,

- 11 65) Cy-C₁₋₁₀ alkyl,
12 66) Cy-C₁₋₁₀ alkenyl, or
13 67) Cy-C₁₋₁₀ alkynyl;

14 L' is a hydrocarbon linker moiety having 1-5 carbon chain atoms and is

15 (i) optionally interrupted by, or terminally attached to, one or more of the following
16 groups:

- 17 1) -C(O)-,
18 2) -O-C(O)-,
19 3) -C(O)-O-,
20 4) -C(O)-NR^c-,
21 5) -NR^c-C(O)-,
22 6) -NR^c-C(O)-NR^d-,
23 7) -NR^c-C(O)-O-,
24 8) -O-C(O)-NR^c-,
25 9) -S(O)_m-,
26 10) -SO₂-NR^c-,
27 11) -NR^c-SO₂-,
28 12) -NR^c-C(NR^m)-,
29 13) -O-,
30 14) -NR^c-, or
31 15) -Cy; or


32 (ii) optionally substituted with one or more substituents independently selected from R^b;

33 L is a hydrocarbon linker moiety having 1-14 carbon chain atoms and is

34 (i) optionally interrupted by, or terminally attached to, one or more of the following
35 groups:

- 36 1) -C(O)-,
37 2) -O-C(O)-,
38 3) -C(O)-O-,
39 4) -C(O)-NR^c-,
40 5) -NR^c-C(O)-,
41 6) -NR^c-C(O)-NR^d-,

- 42 7) $\text{-NR}^c\text{-C(O)-O-}$,
 43 8) $\text{-O-C(O)-NR}^c\text{-}$,
 44 9) $\text{-S(O)}_m\text{-}$,
 45 10) $\text{-SO}_2\text{-NR}^c\text{-}$,
 46 11) $\text{-NR}^c\text{-SO}_2\text{-}$,
 47 12) -O- ,
 48 13) $\text{-NR}^c\text{-}$, or
 49 14) Cy; or
 50 (ii) optionally substituted with one or more substituents independently selected from R^b ;
 51 and
 52 R^3 is alkyl, alkenyl, alkynyl, cycloalkyl, aryl-fused cycloalkyl, cycloalkenyl, aryl, aralkyl,
 53 aryl-substituted alkenyl or alkynyl, cycloalkyl-substituted alkyl, cycloalkenyl-substituted
 54 cycloalkyl, biaryl, alkenoxy, alkynoxy, aralkoxy, aryl-substituted alkenoxy or alkynoxy,
 55 alkylamino, alkenylamino or alkynylamino, aryl-substituted alkylamino, aryl-substituted
 56 alkenylamino or alkynylamino, aryloxy, arylamino, heterocyclyl, heterocyclyl-substituted
 57 alkyl, heterocyclyl-substituted amino, carboxyalkyl substituted aralkyl, or oxocarbocyclyl-
 58 fused aryl; or a moiety of the following formula:



- 59
 60 wherein:
 61 Y^5 is selected from the group consisting of -CO- , -O-CO- , $\text{-SO}_2\text{-}$ and $\text{-PO}_2\text{-}$;
 62 each of R^4 and R^6 , independently, is alkyl, alkenyl, alkynyl, cycloalkyl, aryl-fused
 63 cycloalkyl, cycloalkenyl, aryl, aralkyl, aryl-substituted alkenyl or alkynyl, cycloalkyl-
 64 substituted alkyl, cycloalkenyl-substituted cycloalkyl, biaryl, alkenoxy, alkynoxy, aralkoxy,
 65 aryl-substituted alkenoxy or alkynoxy, alkylamino, alkenylamino or alkynylamino, aryl-
 66 substituted alkylamino, aryl-substituted alkenylamino or alkynylamino, aryloxy, arylamino,
 67 heterocyclyl, heterocyclyl-substituted alkyl, heterocyclyl-substituted amino, carboxyalkyl
 68 substituted aralkyl, oxocarbocyclyl-fused aryl, or an amino acid side chain selected from the
 69 group consisting of arginine, asparagine, glutamine, S-methyl cysteine, methionine and

- 70 corresponding sulfoxide and sulfone derivatives thereof, cyclohexylalanine, leucine,
71 isoleucine, allo-isoleucine, tert-leucine, norleucine, phenylalanine, phenylglycine, tyrosine,
72 tryptophan, proline, alanine, ornithine, histidine, glutamine, norvaline, valine, threonine,
73 serine, beta-cyanoalanine, 2-aminobutyric acid and allothreonine; and
74 R^5 is hydrogen, aryl, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, or aryl-substituted
75 alkyl, R^5 and R^6 may be taken together with the atoms to which they are attached to form a
76 heterocycle of 5 to 7 members;
- 77 each of said Cy is cycloalkyl, heterocyclyl, aryl, or heteroaryl;
- 78 each of said alkyl, alkenyl and alkynyl is optionally substituted with one to four
79 substituents independently selected from R^a ; and
- 80 each of said cycloalkyl, cycloalkenyl, heterocyclyl, aryl, or heteroaryl is optionally
81 substituted with one to four substituents independently selected from R^b ;
- 82 R^a is selected from the group consisting of
- 83 1) Cy,
 - 84 2) $-OR^c$,
 - 85 3) $-NO_2$,
 - 86 4) -halogen,
 - 87 5) $-S(O)_mR^c$,
 - 88 6) $-SR^c$,
 - 89 7) $-S(O)_2OR^c$,
 - 90 8) $-S(O)_2NR^cR^d$,
 - 91 9) $-NR^cR^d$,
 - 92 10) $-O(CR^cR^f)_nNR^cR^d$,
 - 93 11) $-C(O)R^d$,
 - 94 12) $-CO_2R^c$,
 - 95 13) $-P(O)(OR^c)(OR^d)$,
 - 96 14) $-P(O)(R^c)(OR^d)$,
 - 97 15) $-S(O)_mOR^c$,
 - 98 16) $-C(O)NR^cR^j$,
 - 99 17) $-CO_2(CR^cR^f)_nCONR^cR^d$,
 - 100 18) $-OC(O)R^c$,

- 101 19) -CN,
102 20) -NR^cC(O)R^d,
103 21) -OC(O)NR^cR^d,
104 22) -NR^cC(O)OR^d,
105 23) -NR^cC(O)NR^dR^e,
106 24) -CR^c(NOR^d),
107 25) -CF₃,
108 26) -OCF₃, or
109 27) oxo

110 wherein Cy is optionally substituted with one to four substituents independently selected
111 from R^b,

112 R^b is

- 113 1) a group selected from R^a,
114 2) C₁₋₁₀ alkyl,
115 3) C₂₋₁₀ alkenyl,
116 4) C₂₋₁₀ alkynyl,
117 5) aryl-C₁₋₁₀ alkyl, or
118 6) heteroaryl-C₁₋₁₀ alkyl,

119 wherein each of alkyl, alkenyl, alkynyl, aryl, and heteroaryl is optionally substituted with a
120 group independently selected from R^g

121 each of R^c, R^d, R^e, and R^f, independently, is

- 122 1) H,
123 2) C₁₋₁₀ alkyl,
124 3) C₂₋₁₀ alkenyl,
125 4) C₂₋₁₀ alkynyl,
126 5) Cy, or
127 6) Cy-C₁₋₁₀ alkyl;

128 wherein each of alkyl, alkenyl, alkynyl and Cy is optionally substituted with one to four
129 substituents independently selected from R^g;

130 R^g is

- 131 1) halogen,

- 132 2) amino,
- 133 3) carboxy,
- 134 4) -COO-C₁₋₄ alkyl,
- 135 5) -P(O)(OH)₂,
- 136 6) -P(O)(OH)(O-C₁₋₄ alkyl),
- 137 7) -P(O)(C₁₋₄ alkyl)₂,
- 138 8) -P(O)(OH)(C₁₋₄ alkyl),
- 139 9) -P(O)(O-C₁₋₄ alkyl)(C₁₋₄ alkyl),
- 140 10) -SO₂-C₁₋₄ alkyl,
- 141 11) -CO-NH₂,
- 142 12) -CO-NH(C₁₋₄ alkyl),
- 143 13) -CO-N(C₁₋₄ alkyl)₂,
- 144 14) C₁₋₄ alkyl,
- 145 15) C₁₋₄ alkoxy,
- 146 16) aryl,
- 147 17) aryl-C₁₋₄ alkoxy,
- 148 18) hydroxy,
- 149 19) CF₃, or
- 150 20) aryloxy;

151 R^m is

- 152 1) H,
- 153 2) C₁₋₁₀ alkyl,
- 154 3) C₂₋₁₀ alkenyl,
- 155 4) C₂₋₁₀ alkynyl,
- 156 5) Cy,
- 157 6) Cy-C₁₋₁₀ alkyl,
- 158 7) C₁₋₁₀ acyl,
- 159 8) cyano,
- 160 9) C₁₋₁₀ alkyl-sulfonyl, or
- 161 10) C₁₋₁₀ alkoxy; and

162 R^j is

- 163 1) H,
164 2) C₁₋₁₀ alkyl,
165 3) C₂₋₁₀ alkenyl,
166 4) C₂₋₁₀ alkynyl,
167 5) cyano,
168 6) aryl,
169 7) aryl-C₁₋₁₀ alkyl,
170 8) heteroaryl,
171 9) heteroaryl-C₁₋₁₀ alkyl, or
172 10) -SO₂R^k,

173 where R^k is C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl, and aryl;

174 R^e and R^d taken together with the atoms to which they are attached optionally form a
175 heterocyclic ring of 5 to 7 members, said ring containing 0-2 additional heteroatoms
176 independently selected from O, N and S;

177 R^e and R^f taken together with the atoms to which they are attached optionally form a
178 ring of 5 to 7 members, said ring containing 0-2 additional heteroatoms independently
179 selected from O, S and N;

180 m is 0, 1, or 2;

181 n is an integer from 1 to 10;

182 provided that when L is saturated and has 1-4 carbon chain atoms,

- 183 (i) L must contain a heteroatom selected from O, S, and N; or
184 (ii) R³ must contain the moiety o-methylphenyl-ureido-phenyl-CH₂-; or
185 (iii) R¹ must contain only one Cy group;
186 or a pharmaceutically acceptable salt thereof.

1 64. The method of claim 63, wherein said compound is compound nos. 5192, 5283, 6696,
2 6697, 6714, 7234, 7256, 7578, 7662, 8221, 8308, 8309, 8341, 8342, 8343, 8367, 8368, 8469,
3 8491, 8554, 8555, 8571, 8642, 8646, 8685, 8689, 8690, 8698, 8749, 8758, 8796, 8797, 8809,
4 9120, 9169, 9171, 9182, 9227, 9264, 9271, 9315, 9418, 9621, 7083, 7200, 7328, 7399, 7788,
5 7855, 8205, 8290, 8291, 8294, 8295, 8304, 8557, 8582, 8583, 8585, 8586, 8606, 8607, 8628,
6 8674, 8684, 8723, 8746, 8929, 9273, or 9275.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/22285

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : Please See Extra Sheet.

US CL : 514/ 326, 422; 546/208: 548/566, 567, 570

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/ 326, 422; 546/208: 548/566, 567, 570

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CAS--structure

EAST/West--vla4

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO99/06432 A (ATHENA NEUROSCIENCE, INC.) 11 February 1999 (11.02.99), see whole article, especially compounds delineated at p.114-123, particularly those on page 122.	60
Y	WO 99/06434 A (ATHENA NEUROSCIENCES, INC.) 11 February 1999 (11.02.99), see whole article, especially compounds delineated on pages 22-36.	60

☐ Further documents are listed in the continuation of Box C.
 ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*G* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

16 OCTOBER 2000

Date of mailing of the international search report

15 NOV 2000

 Name and mailing address of the ISA/US
 Commissioner of Patents and Trademarks
 Box PCT
 Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

CELIA CHANG

Telephone No. (703) 308-1235

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/22285

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☒ Claims Nos.: 1-59, 61-64
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

Please See Extra Sheet.

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/22285

A. CLASSIFICATION OF SUBJECT MATTER:

IPC (7):

IPC7 A61K 31/40, 31/4025, 31/445; C07D 401/06, 08, 10, 12

BOX I. OBSERVATIONS WHERE CLAIMS WERE FOUND UNSEARCHABLE

2. Where no meaningful search could be carried out, specifically:

Claims 1-59 and 61-64 described the invention in such confusing and ambiguous manner with enormous number of permutation of the parameters with one substituted by another and vice versa i.e. Ra substituted by Rb. Rb can also be substituted by Ra etc., thus, no meaningful search can be conducted. The compounds of pages 18 on are confusing since it is unclear whether each line item corresponding to two different Markush formula are two compounds based on the wherever parameter applies or are one compound. thus, no meaningful search can be made with respect to claims with such line numbers.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☒ **FADED TEXT OR DRAWING**
- ☒ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.